

THE ESI/KECK II DAMPED Ly α ABUNDANCE DATABASE

JASON X. PROCHASKA^{1,2}, ERIC GAWISER^{1,3,4,5}, ARTHUR M. WOLFE^{1,3}, JEFF COOKE^{1,3}, & DAWN GELINO^{1,3}

Accepted to the Astrophysical Journal Supplements: April 10, 2003

ABSTRACT

This paper presents chemical abundance measurements for 37 damped Ly α systems at $z > 2.5$ observed with the Echelle Spectrograph and Imager on the Keck II telescope. We measure the HI column densities of these damped systems through Voigt profile fits to their Ly α profiles and we implement the apparent optical depth method to determine ionic column densities. Figures and tables of all relevant data are presented. A full analysis of the chemical enrichment history described by these observations will be presented in a future paper. This dataset is also valuable for efficiently planning future echelle observations and for rough abundance pattern analyses. We aim to make this entire data set public within three years of this publication.

Subject headings: galaxies: abundances — galaxies: chemical evolution — quasars : absorption lines — nucleosynthesis

1. INTRODUCTION

For nearly a decade our group has been pursuing chemical abundance measurements of the damped Ly α systems (DLA) through high resolution spectroscopy using the HIRES instrument (Vogt et al. 1994) on the Keck I telescope (Prochaska & Wolfe 1999; Prochaska et al. 2001). With over 20 nights of observing, we have built a sample of $N \approx 40$ damped Ly α systems at redshift $z = 1.7$ to 4.5. This dataset has provided a measure of the enrichment history at $z > 2$ (Prochaska & Wolfe 2000), examined nucleosynthetic processes and dust depletion in young protogalaxies (Prochaska & Wolfe 2002), and is now enabling measurements of the star formation rate and history of these galaxies (Prochaska et al. 2002; Wolfe, Prochaska, & Gawiser 2003). A number of groups are currently pursuing similar programs with new echelle instruments on 10-m class telescopes (Molaro et al. 2000; Petitjean et al. 2000; Ellison, Ryan, & Prochaska 2001; Dessauges-Zavadsky et al. 2001; Lopez et al. 2002; Pettini et al. 2002). Progress is limited, however, by the expense of echelle observations; typical high- z quasars require an entire night of observation for reliable analysis of their absorption systems.

On August 29, 1999, the Echelle Spectrograph and Imager (ESI; Sheinis et al. 2002) saw first light on the Keck II telescope. Shortly after, we turned to ESI to perform a new survey of damped Ly α chemical abundances (Prochaska, Gawiser, & Wolfe 2001). Our observing strategy took advantage of ESI's unique characteristics – high throughput, complete wavelength coverage from $\lambda \approx 4000$ –11000Å, both moderate ($R \approx 10000$) and low resolution modes – to address several key issues related to chemical evolution (e.g. Prochaska & Wolfe 2000). In particular, we

focused on $z > 2.5$ damped systems where fewer echelle observations had been made. In four combined nights of integration, we have acquired abundance measurements for 37 damped Ly α systems. In comparison with our HIRES observations, then, we have observed the same number of damped systems in $\approx 1/5$ the observing time. Although the lower resolution of ESI limits the scientific impact of the dataset, the results from this survey will significantly improve our understanding of the chemical evolution history of the early universe (Prochaska et al. 2003, in preparation). The observations also provide valuable 'first-look' analysis crucial to efficiently guide follow-up observations with echelle spectroscopy (e.g. Prochaska, Howk, & Wolfe 2003). Furthermore, our observations include ≈ 10 sub-DLA with $N(\text{HI}) \approx 10^{19-20.3} \text{ cm}^{-2}$ (e.g. Peroux et al. 2001) and unparalleled spectral coverage at $\lambda > 8000\text{Å}$ valuable for both DLA and $z \approx 2$ Mg II metal-line studies.

In this paper, we present the chemical abundance measurements for our ESI damped Ly α sample. The goal of this paper is to present the data and ionic column densities for the damped systems with little accompanying analysis. Future papers will examine the implications of this dataset and extend the analysis to the sub-DLA and $z \approx 2$ Mg II systems. This paper is outlined as follows. We describe the observations and data reduction routines in § 2. In § 3 we detail the methods used to perform the abundance measurements and detail specific issues related to the ESI instrument. The individual damped Ly α systems are presented in § 4 with full figures and tables and we present a brief summary in § 5.

2. OBSERVATIONS AND DATA REDUCTION

This observing program totaled 5 scheduled nights which netted ≈ 4 total nights of scientific observations. In Table 1 we present a comprehensive journal of observations for the quasars included in this paper. Column 1 gives the name, column 2 and 3 list the RA and DEC (J2000), column 4 is the apparent magnitude of the quasar (R for $z > 3$, V for $z < 3$), column 5 is the emission redshift, column 6 lists z_{abs} for the DLA along the sightline, column 7 gives the date of the observation, column 8 details the mode of observation (LWD is the low-dispersion prism mode; ECH is the echelle mode), column 9 shows the slit width (predominantly $0.5''$), column 10 is the exposure

¹ Visiting Astronomer, W.M. Keck Telescope. The Keck Observatory is a joint facility of the University of California, the California Institute of Technology, and NASA.

² UCO/Lick Observatory, University of California, Santa Cruz, Santa Cruz, CA 95064

³ Department of Physics, and Center for Astrophysics and Space Sciences, University of California, San Diego, C-0424, La Jolla, CA 92093-0424

⁴ NSF Postdoctoral Fellow, Yale University, New Haven, CT, PO Box 208101, New Haven, CT 06520

⁵ Andes Prize Fellow, Universidad de Chile, Casilla 36-D, Santiago, Chile

TABLE 1
JOURNAL OF OBSERVATIONS

Quasar	RA (2000)	DEC (2000)	V/R	z_{em}	z_{abs}	Date	Mode	Slit	Exp	Ref
SDSSp0127-00	01 27 00.7	−00 45 59	18.4	4.06	3.72	22jan01	ECH	0.75''	1800	1
						19jul01	ECH	0.50''	1800	
PSS0133+0400	01 33 40.4	+04 00 59	17.9	4.13	3.69,3.77	22jan01	ECH	0.75''	1800	2
PSS0134+3317	01 34 21.6	+33 07 56	18.9	4.52	3.76	15oct01	ECH	0.5''	6000	2,3
PSS0209+0517	02 09 44.7	+05 17 14	17.4	4.18	3.66,3.86	22jan01	ECH	0.75''	1200	2
BRJ0426-2202	04 26 10.3	−22 02 17	18.0	4.30	2.98	22jan01	ECH	0.5''	1800	3,4
FJ0747+2739	07 47 11.2	+27 39 04	17.5	4.11	3.42,3.90	15oct01	ECH	0.5''	6000	5
PSS0808+5215	08 08 49.5	+52 15 16	18.8	4.45	3.11	07apr00	LWD	1.0''	400	2,6
						07apr00	ECH	0.75''	2400	
FJ0812+32	08 12 40.8	+32 08 08	17.6	2.70	2.07,2.63	22jan01	ECH	0.5''	1200	5
Q0930+28	09 33 37.8	+28 45 35	17.5	3.42	3.23	22jan01	ECH	0.5''	1200	7
PC0953+4749	09 56 25.2	+47 34 44	19.5	4.46	3.40,3.89,4.24	22jan01	ECH	0.5''	4800	8,9
PSS0957+3308	09 57 44.5	+33 08 24	17.6	4.25	3.28,4.18	07apr00	LWD	1.0''	400	2,6
						07apr00	ECH	0.75''	1200	
BQ1021+3001	10 21 56.8	+30 01 31	17.0	3.12	2.95	22jan01	ECH	0.75''	1800	5,10
CTQ460	10 39 09.4	−23 13 26	18.0	3.13	2.77	22jan01	ECH	0.5''	1800	11,12
HS1132+2243	11 35 08.0	+22 27 07	17.4	2.88	2.78	22jan01	ECH	0.5''	1200	13
Q1209+0919	12 11 34.1	+09 02 17	18.5	3.30	2.58	22jan01	ECH	0.5''	1800	14
PSS1248+3110	12 48 20.2	+31 10 44	18.9	4.35	3.696	07apr00	LWD	1.0''	400	2,6
						07apr00	ECH	0.75''	2400	
PSS1253-0228	12 53 36.3	−02 28 08	18.7	4.01	2.78	22jan01	ECH	0.5''	2700	2
Q1337+11	13 40 02.4	+11 06 30	19.0	2.92	2.79	22jan01	ECH	0.5''	2400	14, 15
PKS1354-17	13 57 05.9	−17 44 06	18.5	3.15	2.78	19jul01	ECH	0.5''	1800	16
PSS1432+3940	14 32 24.9	+39 40 24	18.6	4.28	3.272	07apr00	LWD	1.0''	400	2,6
						07apr00	ECH	0.75''	3600	
Q1502+4837	15 02 27.3	+48 37 09	17.9	3.20	2.56	19jul01	LWD	1.0''	300	17
						19jul01	ECH	0.50''	1800	
PSS1506+5220	15 06 54.6	+52 20 05	18.1	4.18	3.22	22jan01	ECH	0.5''	1200	2
PSS1723+2243	17 23 23.2	+22 43 58	18.2	4.52	3.69	19jul01	ECH	0.5''	1800	2
PSS2155+1358	21 55 02.1	+13 58 26	17.9	4.26	3.31	31oct00	ECH	0.75''	1800	2
Q2223+20	22 25 37.0	+20 40 18		3.56	3.12	15oct01	ECH	0.5''	3000	9,18
PSS2241+1352	22 41 47.9	+13 52 03	18.7	4.44	4.28	19jul01	ECH	0.50''	2700	2
PSS2323+2758	23 23 41.0	+27 58 01	18.5	4.18	3.68	19jul01	ECH	0.50''	3600	2
FJ2334-09	23 34 46.4	−09 08 12	17.9	3.33	3.06	15oct01	ECH	0.5''	2400	5
Q2342+34	23 44 51.1	+34 33 47	18.6	3.01	2.91	19jul01	ECH	0.50''	1800	18,19
PSS2344+0342	23 44 03.2	+03 42 26	18.6	4.30	3.22	19jul01	ECH	0.50''	1800	2,3

Key to References – 1: Fan et al. (2001); 2: Djorgovski et al. (1998); 3: Peroux et al. (2001); 4: Storrie-Lombardi et al. (2001); 5: White et al. (2000); 6: Prochaska, Gawiser, & Wolfe (2001); 7: Lu et al. (1998); 8: Schneider et al. (1991); 9: Storrie-Lombardi and Wolfe (2000); 10: Gregg et al. (1996); 11: Maza et al. (1995); 12: Lopez et al. (2001); 13: Hagen et al. (1999); 14: Wolfe et al. (1995); 15: Sargent, Steidel & Boksenberg (1989); 16: Schilizzi & McAdam (1975); 17: Snellen et al. (2001); 18: Griffith et al. (1990); 19: White, Kinney, & Becker (1993)

time, and column 11 provides comments.

With this program as the impetus, Prochaska has built an IDL data reduction package for the ESI instrument. Because the ESI modes are nearly static (i.e. there is only one setup per mode) and flexure is well compensated by an active control system, we found it advantageous to create a package tailored to ESI's specific characteristics. The general approach for spectroscopy (the package also reduces images taken with ESI) is similar to the packages designed by the Sloan Digital Sky Survey (SDSS) and Deep Extragalactic Evolutionary Probe (DEEP) teams for their spectroscopic observations. The ESI package constructs a 2D wavelength map in the original data frame from the arc calibration images and implements 1D BSPLINE algorithms for sky subtraction and standard approaches for extraction and coadding. The package is publically available⁶ and will be intermittently supported for the duration of ESI. We now present a more detailed description of the code for ECH mode reductions. Future papers will elaborate on the reduction routines for the LWD and imaging modes.

At the heart of the code is an IDL structure designed to organize all data from a single night of observing. This structure contains the vital characteristics of a given frame (e.g. ESI mode, exposure time, slit size) and helps guide the reduction process. In general, one passes this structure to a given IDL routine and the code automatically coordinates and processes the frames. Bias subtraction is performed using a bias image constructed from a series of 0s exposures. This approach is advantageous to using the overscan because of a significant bias variation in the sector of the CCD containing two chip defects. A normalized flat is constructed (ideally from dome flats as the internal quartz lamps exhibit significant scattered light) to remove pixel-to-pixel variations. In the process, the slit-edges of each echellette order are determined from the flat and if the data were taken in the standard two-amp mode, then the gain ratio is calculated. This gain ratio is applied to all data taken throughout the night which is effective but not perfect. In particular, data near 5570Å (the wavelength affected by the boundary between amplifiers) may suffer if the electronics are not perfectly stable throughout the night.

The most important step in the ECH routine is the construction of a 2D wavelength image which contains a unique wavelength value for every data pixel in the ECH image. To create this wavelength image, the code first processes and combines all arc frames for a given slit size. In sequence, it straightens⁷ each echellette order and extracts the central few columns (the data runs parallel to the columns in ESI) to provide a 1D arc spectrum. This spectrum is cross-correlated against an archived 1D spectrum to determine any integer offset. We have found that ESI is stable to better than 3 pixels for data taken over the two years since commissioning. After applying this offset, the program automatically identifies arc lines assuming the wavelength solution of the archived 1D spectrum. It then centroids these lines and determines a final 1D solution

by fitting a polynomial with order between 5 and 9. The code traces bright, individual arc lines to determine their curvature relative to the slit which is typically ≈ 0.1 pix from slit-edge to slit-edge. A 2D polynomial is simultaneously fit to the centroids of all of the arc lines of a given echellette order and a 2D wavelength map is created with typical RMS < 2 km/s. Finally, this wavelength map is inverted to the original data frame to create a 2D wavelength image (vacuum corrected) that is used by all data frames with identical slit width. For a given exposure, every data pixel now has a unique λ , flux pair.

Sky subtraction, extraction and the creation of a final 1D spectrum proceed in relatively standard fashion. The data images are bias subtracted and flattened. All significant objects are identified within the slit and masked for sky subtraction. For the bluest orders where sky lines are negligible, a low order polynomial is fit row by row to the sky and subtracted from the data. For the orders with $\lambda > 5000$ Å, the sky subtraction algorithm utilizes the 2D wavelength image. The λ , flux pairs for all sky pixels in a given order are collapsed to 1D and fit with a 1D BSPLINE to create a single, functional representation of the sky at all wavelength. This approach is particularly advantageous for fitting bright sky lines and the user can modify the location of the BSPLINE breakpoints to improve the fitting of specific sky lines as desired. The 1D BSPLINE solution is evaluated at every data pixel and subtracted from the image. The principal science object is extracted from the sky-subtracted image using a boxcar aperture (determined uniquely for each echellette order) and the λ , flux pairs are collapsed for each order to produce ten 1D spectra with 10 km/s pixels. Note that during this process the λ values are heliocentric corrected. The error array is derived in a similar fashion and includes uncertainty due to object, sky, and readnoise. Cosmic rays are identified during the extraction process by comparing the collapsed object profile to the row-by-row profile. These are then flagged and all data at that wavelength are rejected from future analysis. At this stage, individual exposures are optimally coadded with rejection and the resulting data is fluxed using a standard star from that night or the archive. Finally, the individual orders are coadded to create a single, continuous 1D spectrum.

Ideally, the package is fully automated with minimal user interaction beyond bookkeeping and optional checking. Several procedures are computationally expensive (construction of the arc image; sky subtraction) in part owing to the number of pixels involved and in part due to the implementation of IDL instead of C or FORTRAN. The current code will reduce and extract a single data frame in ≈ 15 min on a 900MHz Sunblade processor. Future generations of the code will include procedures for optimal extraction and generic binning modes and will also include updated archival calibration frames.

3. METHODS

The techniques employed in this paper to determine column densities and errors do not differ significantly from those of our previous work with echelle data (e.g. Prochaska et al. 2001). In general, we implement the apparent optical depth method (AODM, Savage and Sembach 1991) to evaluate column densities of unsaturated, unblended

⁶ <http://www2.keck.hawaii.edu/realpublic/inst/esi/ESIRedux/index.html>

⁷ The curvature of the echellette orders is determined from a sequence of pin-hole observations taken either by the observer or from the calibration archive.

metal-line transitions. We stress one difference from our previous work: we establish a stricter saturation limit to prevent “hidden saturation” allowed by the lower resolution of the ESI instrument. Also note that the upper limits reported are formally 3σ statistical limits and the measurement uncertainties for detections are reported as 1σ errors derived from the AODM. Unless otherwise noted, these limits were derived by integrating the optical depth over the velocity range of the spectra covering all of the absorption in the detected transitions. Because these uncertainties do not take into account systematic error (i.e. continuum placement), we caution the reader when interpreting them. For example, a reported error of 0.01 dex is too optimistic for this dataset. We recommend adopting a minimum error of 0.1 dex for the measurements of single transitions. In turn, we would recommend assuming that relative abundances (e.g. Si/Fe) have at least 0.15 dex uncertainty. Finally, the incidence of contradiction between our reported limits and the measured value for ions with multiple observed transitions suggests the limits refer to $\approx 95\%$ c.l.




FIG. 1.— Observed differences δ between column density measurements as a function of observed equivalent width (EW) for a series of metal-line transitions from three DLA (medium gray squares: PH957; light gray circles: FJ0812+32; dark gray stars: PSS0957+33, $z = 4.18$) observed with both ESI and HIRES. Because of the effects of resolution, the ESI values are significantly lower when $EW > 1\text{\AA}$. There is a milder effect for transitions with $0.6 < EW < 1\text{\AA}$.

The accuracy of our ESI measurements and the effects of saturation are well demonstrated in Figure 1. In the figure, we plot $\delta_N \equiv \log N(X)_{ESI} - \log N(X)_{HIRES}$ against $EW_{ESI}(X)$, the observed equivalent width of transition X for a set of transitions in three damped Ly α systems with both ESI and HIRES observations. For transitions with $EW < 0.6\text{\AA}$, the differences are consistent with measurement error while transitions with $EW > 1\text{\AA}$ clearly suffer from saturation in the ESI observations. For transitions with $0.6 < EW < 1\text{\AA}$, the ESI observations are systematically low but the effect is mild. In the following analysis, we impose lower limits to the column densities measured from transitions with EW exceeding $\approx 0.6\text{\AA}$.




FIG. 2.— A comparison of C^{3+} column densities measured independently from the C IV 1548 and 1550 transitions in the same DLA using the apparent optical depth method (AODM). The dashed line traces the line of equality. The good agreement between the two values over a large range in $N(C^{3+})$ indicates the AODM is an accurate method for measuring column densities in unblended transitions.

In Figure 2 we present a quantitative test of our error analysis by plotting $N(C^{3+})$ measured independently from the C IV 1548 and 1550 transitions of ≈ 20 damped systems. With the exception of one or two data points which may suffer from unidentified blending, the measurements are consistent with a reduced $\chi^2_\nu \approx 1$. One notes no obvious effect from line saturation although several obvious cases are not included in Figure 2.

To determine the $N(\text{HI})$ value of the damped Ly α systems in our sample, we have analysed the fluxed Ly α profiles from our echellette observations. We have found this to be a crucial aspect of the Ly α profile fitting process. In particular, we suspect that by continuum fitting the data prior to performing the Ly α profile analysis that we tended to underestimate the $N(\text{HI})$ values. It is possible that this systematic error holds for other studies where the authors continuum fit the quasar prior to the damped Ly α analysis. Although the absolute fluxing of the ESI data is not accurate because of uncorrected slit losses, the relative fluxing is far superior to results obtained for HIRES and other echelle observations. In short, we have found that the resolution and fluxing of the ESI instrument is nearly ideal for this analysis although we suspect that higher sampling would be advantageous. We emphasize that our analysis is not strictly quantitative; the adopted $N(\text{HI})$ values do not correspond to a minimization of χ^2 for regions of the spectrum. Instead, we present our favored value and place conservative error estimates to these values. Our experience with line-fitting is that standard χ^2 analysis would yield unrealistically small error estimates, in particular because the continuum uncertainty is difficult to quantify. The error estimates provided in this paper are sufficiently conservative, however, that the reader should consider them to be roughly 95% c.l.

To determine $N(\text{HI})$ we tied the centroid of a Voigt pro-

TABLE 2
ATOMIC DATA

Transition	λ	f	Ref
HI 972	972.5368	0.02900000	1
OI 988	988.7734	0.04318000	1
NIII 989	989.7990	0.10660000	1
+ HI 1025	1025.7223	0.07912000	1
CII 1036	1036.3367	0.12310000	1
OI 1039	1039.2304	0.00919700	1
NII 1083	1083.9900	0.10310000	1
FeIII 1122	1122.5260	0.05390000	16
NI 1134	1134.1653	0.01342000	1
NI 1134	1134.4149	0.02683000	1
NI 1134	1134.9803	0.04023000	1
FeII 1144	1144.9379	0.10600000	2
SiII 1190	1190.4158	0.25020000	1
SiII 1193	1193.2897	0.49910000	1
NI 1199	1199.5496	0.13000000	1
NI 1200	1200.2233	0.08620000	1
NI 1200	1200.7098	0.04300000	1
SiIII 1206	1206.5000	1.66000000	1
HI 1215	1215.6701	0.41640000	1
MgII 1239	1239.9253	0.00063300	3
MgII 1240	1240.3947	0.00035500	3
SII 1250	1250.5840	0.00545300	1
SII 1253	1253.8110	0.01088000	1
SII 1259	1259.5190	0.01624000	1
SiII 1260	1260.4221	1.00700000	1
CI 1277	1277.2450	0.09665000	1
OI 1302	1302.1685	0.04887000	1
SiII 1304	1304.3702	0.09400000	4
NiII 1317	1317.2170	0.07786000	1
CII 1334	1334.5323	0.12780000	1
CII* 1335	1335.7077	0.11490000	1
OI 1355	1355.5977	0.00000124	1
NiII 1370	1370.1310	0.07690000	5
SiIV 1393	1393.7550	0.52800000	1
SiIV 1402	1402.7700	0.26200000	1
NiII 1454	1454.8420	0.03230000	7
SiII 1526	1526.7066	0.12700000	9
CIV 1548	1548.1950	0.19080000	1
CIV 1550	1550.7700	0.09522000	1
CI 1560	1560.3092	0.08041000	1
FeII 1608	1608.4511	0.05800000	17
FeII 1611	1611.2005	0.00136000	16
CI 1656	1656.9283	0.14050000	1
AlII 1670	1670.7874	1.88000000	1
NiII 1703	1703.4050	0.00600000	7
NiII 1709	1709.6042	0.03240000	7
NiII 1741	1741.5531	0.04270000	7
NiII 1751	1751.9157	0.02770000	7
SiII 1808	1808.0130	0.00218600	11
AlIII 1854	1854.7164	0.53900000	1
AlIII 1862	1862.7895	0.26800000	1
TiII 1910a	1910.6000	0.20200000	12
TiII 1910b	1910.9380	0.09800000	12
ZnII 2026	2026.1360	0.48900000	13
CrII 2056	2056.2539	0.10500000	13
CrII 2062	2062.2340	0.07800000	13
ZnII 2062	2062.6640	0.25600000	13
CrII 2066	2066.1610	0.05150000	13
FeII 2249	2249.8768	0.00182100	14
FeII 2260	2260.7805	0.00244000	14
FeII 2344	2344.2140	0.11400000	3
FeII 2374	2374.4612	0.03130000	3
FeII 2382	2382.7650	0.32000000	3
MnII 2576	2576.8770	0.35080000	1
FeII 2586	2586.6500	0.06910000	3
MnII 2594	2594.4990	0.27100000	1
FeII 2600	2600.1729	0.23900000	3
MgII 2796	2796.3520	0.61230000	15
MgII 2803	2803.5310	0.30540000	15
MgI 2852	2852.9642	1.81000000	1

Key to References – 1: Morton (1991); 2: Howk et al. (2000); 3: Morton (2003); 4: Tripp et al. (1996); 5: Fedchak & Lawler (1999); 7: Fedchak, Wiese, & Lawler (2000); 9: Schectman et al. (1998); 11: Bergeson & Lawler (1993b); 12: Wiese, Fedchak, & Lawler (2001); 13: Bergeson & Lawler (1993); 14: Bergeson, Mullman, & Lawler (1994); 15: Verner (1996); 17: Bergeson et al. (1996)

TABLE 3
SOLAR ABUNDANCES

Elm	ϵ^a
Al	6.49
Ar	6.52
As	2.37
B	2.79
C	8.59
Ca	6.35
Cl	5.28
Co	4.91
Cr	5.67
Cu	4.29
Fe	7.50
Ga	3.13
Ge	3.63
H	12.00
K	5.13
Kr	3.23
Li	3.31
Mg	7.58
Mn	5.53
N	7.93
Na	6.32
Ne	8.08
Ni	6.25
O	8.74
P	5.53
Pb	2.06
S	7.20
Si	7.56
Sn	2.14
Ti	4.94
Xe	2.23
Zn	4.67
Zr	2.61

^aMeteoritic abundances except C,N,O

file to the metal-line transitions⁸ and varied the $N(\text{HI})$ value and the quasar continuum simultaneously to achieve a solution which ideally matched both the core and wings of the profile. In all cases, we required that the solution never fell significantly below the data values in any portion of the Ly α profile. Nevertheless, because unidentified Ly α blending is always present, it is plausible that the resulting $N(\text{HI})$ values are over-estimated in some cases. On the other hand, one is systematically likely to underestimate the quasar continuum and therefore underestimate $N(\text{HI})$. It would be valuable to fully assess these effects with numerical simulations. As is standard practice in DLA searches we rejected Broad Absorption Line quasars which make DLA identification extremely difficult and do not consider systems found within 3000 km/s of the quasar emission redshift to be DLAs owing to the chance that these systems are physically associated with the quasar (but see Møller, Warren, & Fynbo 1998; Ellison et al. 2002).

Throughout the paper, we adopt the wavelengths and oscillator strengths presented in Table 2. When possible, we have adopted laboratory values for the oscillator strengths. We also adopt solar meteoritic abundances from Grevesse et al. (1996) and Holweger (2001) for the elemental abundances listed in the paper (Table 3).

⁸ Even when the metal-line transitions show broad (up to 200 km/s), complex kinematics we are able to estimate the systematic velocity centroid to within ~ 30 km/s i.e. $\delta z \simeq 0.0001$.

4. INDIVIDUAL SYSTEMS

In this section we present the metal-line and Ly α profiles of the DLA and tabulate the measured ionic column densities. Regarding the metal-line figures, $v = 0$ km s⁻¹ is arbitrarily defined and blends from coincident absorption lines are indicated by dotted lines. The normalized continuum is shown by the dotted line and the zero-flux line is marked by a dash-dot line. For the low-ion transitions – the dominant ionic transition of a given element in an HI region – we tabulate N_{adopt} by taking the weighted mean (or limit) of all measured transitions. For these cases, we also convert N_{adopt} into an elemental abundance $[\text{X}/\text{H}]$ assuming no ionization corrections and adopting solar meteoritic abundances. In the Ly α figures, a vertical dashed line indicates the centroid of the Ly α profile while the dotted line traces the quasar continuum. Furthermore, the best fit solution for the Ly α profile is traced by a solid line and is bounded by two dash-dot lines which indicate our estimate of the error in the Ly α solution.




FIG. 3.— Ly α profile of the damped Ly α system at $z = 3.727$ toward SDSS0127-00. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.15^{+0.10}_{-0.10}$. Examining the region at $\lambda \approx 5760\text{\AA}$, one notes the data have flux significantly higher than the overplotted Voigt profile. This is due an incorrect matching in the gain of the two amplifiers for ESI (§ 2). This solution is well constrained by the left wing of the Ly α profile and is simplified by the relatively well behaved continuum (dotted line) across this region.

4.1. SDSS0127-00, $z = 3.727$

The quasar SDSS0127-00 is a member of the Early Data Release of the Sloan Digital Sky Survey (Fan et al. 2001). Even in the SDSS discovery spectrum, a damped Ly α system is obvious at $\approx 5770\text{\AA}$. In Figure 3 we present the Ly α profile and our solution which is well constrained by the left wing of the profile. One notes a complication with the current IDL package: normalizing the gain of the two amplifiers. This issue is only manifest at $\lambda \approx 5770\text{\AA}$ in the spectrum and only arises when the data frame had a significantly different gain ratio from the flats. Of our observations, this is the most severe case. Future versions of




FIG. 4.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.727$ toward SDSS01027-00. The vertical line at $v = 0$ corresponds to $z = 3.72743$.

TABLE 4

IONIC COLUMN DENSITIES: SDSS0127-00, $z = 3.727$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 14.672	> 14.672	> -3.068
C II*	1335.7	13.139 ± 0.064		
C IV	1550.8	14.129 ± 0.036		
N I	1200.2	14.028 ± 0.037	14.028 ± 0.037	-3.052 ± 0.107
O I	1302.2	> 15.185	> 15.185	> -2.705
Al II	1670.8	> 12.743	> 12.743	> -2.897
Si II	1304.4	> 14.306	> 14.306	> -2.404
Si II	1526.7	> 14.221		
Si II	1808.0	< 14.760		
Si IV	1402.8	13.439 ± 0.049		
Ni II	1741.6	< 13.691	< 13.691	< -1.709

the ESI IDL package will provide a means for correcting

this effect. The metal-line profiles of this damped system are shown in Figure 4 and the ionic column densities are listed in Table 4.




FIG. 5.— Ly α profiles of the damped Ly α systems at $z = 3.693$ and 3.774 toward PSS0133+0400. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.70^{+0.10}_{-0.15}$ and $\log N(\text{HI}) = 20.55^{+0.10}_{-0.15}$ respectively. The vertical dash-dot lines correspond to coincident Ly α forest lines that we included in the analysis. While the $z = 3.693$ DLA is well constrained by the observations, the core of the $z = 3.77$ DLA is significantly wider than the Ly α wings suggest. Therefore, this DLA is more sensitive to the continuum placement (dotted line) and the $N(\text{HI})$ value is more uncertain. Note that we introduced additional components to investigate their effect on the DLA profiles and found minimal effect.

TABLE 5

IONIC COLUMN DENSITIES: PSS0133+0400, $z = 3.693$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 14.160	> 14.160	> -3.130
C II*	1335.7	< 12.563		
C IV	1548.2	13.523 ± 0.028		
C IV	1550.8	< 13.787		
Al II	1670.8	12.372 ± 0.031	12.372 ± 0.031	-2.818 ± 0.129
Si II	1526.7	> 14.256	> 14.256	> -2.004
Si II	1808.0	< 14.546		
Si IV	1402.8	< 12.724		
Fe II	1608.5	13.511 ± 0.065	13.511 ± 0.065	-2.689 ± 0.141

4.2. PSS0133+0400, $z = 3.693, 3.774$

We identified two damped systems toward this quasar from the Palomar Sky Survey (PSS Djorgovski et al. 1998). The Ly α profiles of the damped systems are shown in Figure 5. Because of their proximity, we found it necessary to fit the two profiles simultaneously. While the damped system at $z = 3.693$ is well constrained by the observations, the core of the $z = 3.774$ DLA is significantly wider than the damping wings suggest. This indicates blending from coincident Ly α forest lines with $N(\text{HI}) > 10^{15} \text{ cm}^{-2}$ and

TABLE 6

IONIC COLUMN DENSITIES: PSS0133+0400, $z = 3.774$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 15.309	> 15.309	> -1.831
C II*	1335.7	< 13.949		
C IV	1548.2	> 14.306		
C IV	1550.8	14.355 ± 0.016		
Al II	1670.8	> 13.792	> 13.792	> -1.248
Al III	1854.7	13.164 ± 0.027		
Al III	1862.8	13.108 ± 0.054		
Si II	1526.7	> 15.079	15.466 ± 0.035	-0.644 ± 0.130
Si II	1808.0	15.466 ± 0.035		
Si IV	1393.8	13.799 ± 0.011		
Si IV	1402.8	13.894 ± 0.015		
Ti II	1910.6	13.041 ± 0.066	13.041 ± 0.066	-0.449 ± 0.141
Cr II	2056.3	< 13.240	< 13.240	< -0.980
Fe II	1608.5	> 14.868	> 14.868	> -1.182
Fe II	1611.2	< 15.375		
Ni II	1370.1	13.999 ± 0.029	13.978 ± 0.026	-0.822 ± 0.128
Ni II	1741.6	13.899 ± 0.066		
Zn II	2026.1	< 13.099	< 13.099	< -0.121

the net result is that the $N(\text{HI})$ value is more sensitive to continuum placement.

Velocity profiles of the metal-line transitions are presented in Figures 6 and 7 and the measured ionic column densities are listed in Tables 5 and 6. The system at $z = 3.774$ is notable for exhibiting the highest DLA metallicity at $z > 3$ and also a relatively low level of differential depletion ($[\text{Si}/\text{Ni}] \approx 0.2$). Follow-up echelle observations of this sightline may be particularly valuable.

TABLE 7

IONIC COLUMN DENSITIES: PSS0134+3317, $z = 3.761$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	13.282 ± 0.103		
C IV	1550.8	13.237 ± 0.138		
Al II	1670.8	12.612 ± 0.027	12.612 ± 0.027	-2.728 ± 0.080
Si II	1193.3	> 13.889	> 13.889	> -2.521
Si II	1808.0	< 14.681		
Si III	1206.5	> 13.815		

4.3. PSS0134+3317, $z = 3.761$

This quasar is a member of the PSS sample and the damped Ly α systems was discovered by Peroux et al. (2001). The Ly α profile corresponding to this DLA is given in Figure 8. Although the profile is coincident with the O VI peak of PSS0134+3307, the HI column density is well constrained by the observations. This DLA has a low metallicity and the majority of the strongest transitions are lost in the Ly α forest. The few valuable transitions are shown in Figure 9 and provide column density limits for only Si⁺ and Al⁺ (Table 7).

4.4. PSS0209+0517, $z = 3.667, 3.864$

This quasar exhibits two damped Ly α systems at $z = 3.667$ and 3.864 (discovered by Peroux et al. 2001) whose Ly α profiles are presented in Figure 10. The cores of both




FIG. 6.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.693$ toward PSS0133+0400. The vertical line at $v = 0$ corresponds to $z = 3.6926$.

TABLE 8

IONIC COLUMN DENSITIES: PSS0209+0517, $z = 3.667$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1560.3	< 12.910		
C IV	1548.2	13.181 ± 0.049		
O I	1302.2	> 14.774	> 14.774	> -2.416
Al II	1670.8	12.521 ± 0.037	12.520 ± 0.037	-2.420 ± 0.107
Al III	1854.7	< 12.217		
Si II	1260.4	> 14.107	> 14.107	> -1.903
Si II	1526.7	> 13.885		
Si II	1808.0	< 14.455		
Si III	1206.5	> 13.846		
Si IV	1402.8	13.161 ± 0.051		
Cr II	2056.3	< 13.079	< 13.079	< -1.041
Fe II	1608.5	13.635 ± 0.053	13.635 ± 0.053	-2.315 ± 0.113
Ni II	1709.6	< 13.249	< 13.250	< -1.450
Ni II	1751.9	< 13.621		

Ly α profiles are well determined and provide a precise determination of $N(\text{HI})$. The metal-line profiles of the two




FIG. 7.— Velocity plot of the metal-line transitions for the damped $\text{Ly}\alpha$ system at $z = 3.774$ toward PSS0133+0400. The vertical line at $v = 0$ corresponds to $z = 3.77356$.

DLA are presented in Figures 11 and 12 and the ionic column densities are listed in Tables 8, 9.

4.5. *BRJ0426-2202*, $z = 2.983$

This quasar was discovered by Storrie-Lombardi et al. (2001) and Peroux et al. (2001) identified the damped $\text{Ly}\alpha$ system. The $\text{Ly}\alpha$ profile of this large HI DLA is difficult to constrain because of a higher redshift Lyman limit system which is significantly suppressing the quasar flux below $4800\text{\AA}$. Nevertheless, the red wing of the DLA is relatively free from contamination and a reasonable fit to the $N(\text{HI})$ value was achieved (Figure 13). We note that our value is 0.4 dex larger than the value reported by Peroux et al. (2001) which may suggest an uncertainty in $N(\text{HI})$ even larger than the 0.15 dex we adopt.

Because the DLA has a substantially lower redshift than the quasar and a low metallicity, it exhibits only a few

TABLE 9
IONIC COLUMN DENSITIES: PSS0209+0517, $z = 3.864$

Ion	λ	AODM	N_{adopt}	$[X/\text{H}]$
C II	1334.5	> 14.020	> 14.020	> -3.120
C II*	1335.7	< 12.946		
C IV	1548.2	13.311 ± 0.066		
C IV	1550.8	13.373 ± 0.095		
O I	1302.2	> 14.368	> 14.368	> -2.922
Al II	1670.8	12.146 ± 0.048	12.146 ± 0.048	-2.894 ± 0.111
Al III	1854.7	< 12.150		
Si II	1260.4	> 13.819	13.463 ± 0.034	-2.647 ± 0.106
Si II	1304.4	13.542 ± 0.044		
Si II	1526.7	13.396 ± 0.052		
Si IV	1393.8	12.922 ± 0.052		
Si IV	1402.8	12.934 ± 0.099		
Fe II	1608.5	< 13.340	< 13.340	< -2.710
Ni II	1370.1	< 13.123	< 13.123	< -1.677
Ni II	1741.6	< 13.236		




FIG. 8.— Ly α profile of the damped Ly α system at $z = 3.761$ toward PSS0134+3317. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.85^{+0.05}_{-0.10}$. Although this DLA is positioned on the O VI peak of the quasar, the core is well resolved and we consider the $N(\text{HI})$ to be well constrained.

TABLE 10
IONIC COLUMN DENSITIES: BRJ0426-2202, $z = 2.983$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.3	< 13.020		
Al II	1670.8	> 12.931	> 12.931	> -3.059
Al III	1854.7	< 12.196		
Si II	1808.0	< 14.996	< 14.996	< -2.064
Cr II	2056.3	< 12.901	< 12.901	< -2.269
Fe II	2260.8	< 14.887	14.152 ± 0.067	-2.848 ± 0.164
Fe II	2344.2	> 14.240		
Fe II	2374.5	14.152 ± 0.067		
Fe II	2382.8	> 13.625		
Ni II	1709.6	< 13.373	< 13.373	< -2.377
Ni II	1741.6	< 13.512		
Ni II	1751.9	< 13.403		
Zn II	2026.1	< 12.165	< 12.165	< -2.005

metal-line transitions (Figure 14). The resulting ionic column densities are given in Table 10.

TABLE 11
IONIC COLUMN DENSITIES: FJ0747+2739, $z = 3.423$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 15.096	> 15.096	> -2.344
C IV	1548.2	> 14.308		
C IV	1550.8	> 14.403		
Al II	1670.8	> 13.204	> 13.204	> -2.136
Al III	1854.7	13.060 ± 0.031		
Si II	1193.3	> 14.481	> 14.539	> -1.871
Si II	1526.7	> 14.539		
Si II	1808.0	< 14.932		
Fe II	1608.5	> 14.430	> 14.430	> -1.920
Fe II	1611.2	< 14.832		
Fe II	2260.8	< 14.701		
Ni II	1751.9	< 13.271	< 13.272	< -1.828




FIG. 9.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.761$ toward PSS0134+3317. The vertical line at $v = 0$ corresponds to $z = 3.76087$.

TABLE 12
IONIC COLUMN DENSITIES: FJ0747+2739, $z = 3.900$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1036.3	> 14.380	> 14.503	> -2.587
C II	1334.5	> 14.503		
C II*	1335.7	13.259 ± 0.075		
C IV	1548.2	13.715 ± 0.014		
O I	1302.2	> 14.806	> 14.806	> -2.434
Al II	1670.8	12.461 ± 0.025	12.461 ± 0.025	-2.529 ± 0.103
Si II	1304.4	14.087 ± 0.015	14.031 ± 0.011	-2.029 ± 0.101
Si II	1526.7	13.983 ± 0.016		
Si II	1808.0	< 14.423		
S II	1253.8	< 14.361	< 14.361	< -1.339
Fe II	1608.5	< 13.799	< 13.799	< -2.201
Ni II	1317.2	< 13.106	< 13.106	< -1.644
Zn II	2026.1	< 12.398	< 12.398	< -0.772




FIG. 10.— Ly α profiles of the damped Ly α systems at $z = 3.667$ and 3.864 toward PSS0209+0517. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.45^{+0.1}_{-0.1}$ and $\log N(\text{HI}) = 20.55^{+0.1}_{-0.1}$ respectively. In both cases, the core of the Ly α profile is relatively unblended with coincident Ly α lines and provides a good determination of $N(\text{HI})$ even though the continuum (dotted line) shows modest variation across the profiles. Note that the enhancement in the continuum at $\lambda \approx 5680\text{\AA}$ is due to O VI emission by the QSO.

4.6. FJ0747+2739, $z = 3.423, 3.900$

This bright FIRST quasar (White et al. 2000) has two, well separated damped Ly α systems along its sightline. The lower redshift DLA (Figure 15) has a very well determined $N(\text{HI})$ value owing to a simple continuum and unblended core. The DLA at $z = 3.900$, however, has a more uncertain continuum owing to its proximity to the Ly α emission peak (Figure 16).

Figures 17 and 18 present the metal-line profiles for the $z = 3.423$ and $z = 3.900$ DLA respectively. Finally, the ionic column densities of the two DLA are provided in Tables 11 and 12.

TABLE 13

IONIC COLUMN DENSITIES: PSS0808+52, $z = 3.113$

Ion	λ	AODM	N_{adopt}	[X/H]
O I	1302.2	> 14.945	> 14.945	> -2.445
Al III	1862.8	13.425 ± 0.014		
Si II	1304.4	> 14.477	14.599 ± 0.120	-1.611 ± 0.139
Si II	1808.0	14.599 ± 0.120		
Cr II	2056.3	< 12.735	< 12.735	< -1.585
Fe II	2249.9	< 14.626	14.170 ± 0.040	-1.980 ± 0.081
Fe II	2344.2	> 13.966		
Fe II	2374.5	14.170 ± 0.040		
Ni II	1741.6	< 13.238	< 13.238	< -1.662
Zn II	2026.1	< 12.134	< 12.134	< -1.186

4.7. PSS0808+52, $z = 3.113$

We presented a first analysis of this damped Ly α system in Prochaska, Gawiser, & Wolfe (2001). Our re-analysis




FIG. 11.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.667$ toward PSS0209+0517. The vertical line at $v = 0$ corresponds to $z = 3.66674$.

yields very similar results. The Ly α profile is shown in Figure 19 and gives a very precise measure of the HI column density. The complete set of metal-line profiles and ionic column densities are presented in Figure 20 and Table 13.




FIG. 12.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.864$ toward PSS0209+0517. The vertical line at $v = 0$ corresponds to $z = 3.86427$.




FIG. 13.— Ly α profile of the damped Ly α system at $z = 2.983$ toward BRJ0426-2202. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.50^{+0.15}_{-0.15}$. The blue wing of the Ly α profile is severely blended with the higher order Lyman series of a higher redshift Lyman limit system. The red wing is reasonably free from contamination, although the resulting fit is sensitive to the continuum placement. This results in a more uncertain determination of $N(\text{HI})$.




FIG. 14.— Velocity plot of the metal-line transitions for the damped $\text{Ly}\alpha$ system at $z = 2.983$ toward BRJ0426-2202. The vertical line at $v = 0$ corresponds to $z = 2.98308$.




FIG. 15.— $\text{Ly}\alpha$ profile of the damped $\text{Ly}\alpha$ system at $z = 3.423$ toward FJ0747+2739. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.85^{+0.05}_{-0.05}$. Its HI value is particularly well determined because of the simplicity of the continuum and the absence of blending with the core of the profile.




FIG. 16.— $\text{Ly}\alpha$ profile of the damped $\text{Ly}\alpha$ system at $z = 3.900$ toward FJ0747+2739. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.50^{+0.10}_{-0.10}$. This DLA is located near the $\text{Ly}\alpha$ peak of the quasar and the red wing is more difficult to determine. Nevertheless, the blue wing and therefore the HI column density is reasonably well constrained, despite the presence of a lower column-density system in the middle of the blue wing which was considered as part of the fit.




FIG. 17.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.423$ toward FJ0747+2739. The vertical line at $v = 0$ corresponds to $z = 3.4233$.




FIG. 18.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.900$ toward FJ0747+2739. The vertical line at $v = 0$ corresponds to $z = 3.9000$.




FIG. 19.— Ly α profile of the damped Ly α system at $z = 3.113$ toward PSS0808+52. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.65^{+0.07}_{-0.07}$. This DLA has a well sampled core and therefore a very precise measurement of $N(\text{HI})$.




FIG. 20.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.113$ toward PSS0808+52. The vertical line at $v = 0$ corresponds to $z = 3.11318$.

TABLE 14
IONIC COLUMN DENSITIES: FJ0812+32, $z = 2.626$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.9	12.890 ± 0.089		
C II	1334.5	> 15.140	> 15.140	> -2.800
C II*	1335.7	> 14.089		
C IV	1548.2	> 14.496		
C IV	1550.8	14.534 ± 0.008		
N I	1159.8	< 17.606	< 17.606	< 0.326
O I	1302.2	> 15.488	> 15.488	> -2.602
O I	1355.6	< 17.981		
Mg II	1239.9	16.313 ± 0.080	16.324 ± 0.065	-0.606 ± 0.119
Mg II	1240.4	16.349 ± 0.111		
Mg II	2796.4	> 13.839		
Mg II	2803.5	> 14.030		
Al II	1670.8	> 13.659	> 13.659	> -2.181
Al III	1854.7	13.175 ± 0.016		
Al III	1862.8	13.243 ± 0.024		
Si II	1304.4	> 15.052	> 15.782	> -1.128
Si II	1526.7	> 14.884		
Si II	1808.0	> 15.782		
Si IV	1393.8	> 14.025		
Si IV	1402.8	14.059 ± 0.011		
S II	1250.6	> 15.438	> 15.438	> -1.112
S II	1253.8	> 15.284		
S II	1259.5	> 15.263		
Cr II	2056.3	13.364 ± 0.034	13.320 ± 0.032	-1.700 ± 0.105
Cr II	2062.2	13.330 ± 0.041		
Cr II	2066.2	13.187 ± 0.081		
Mn II	2594.5	13.319 ± 0.037	13.319 ± 0.037	-1.561 ± 0.107
Fe II	1125.4	15.121 ± 0.079	15.097 ± 0.025	-1.753 ± 0.103
Fe II	1144.9	> 14.790		
Fe II	1608.5	> 14.743		
Fe II	1611.2	15.132 ± 0.058		
Fe II	2249.9	15.153 ± 0.035		
Fe II	2260.8	15.019 ± 0.046		
Fe II	2344.2	> 14.528		
Fe II	2374.5	> 14.764		
Fe II	2382.8	> 14.303		
Ni II	1317.2	13.825 ± 0.038	13.846 ± 0.019	-1.754 ± 0.102
Ni II	1370.1	13.930 ± 0.034		
Ni II	1454.8	13.754 ± 0.070		
Ni II	1703.4	< 14.003		
Ni II	1741.5	13.850 ± 0.037		
Ni II	1751.9	13.796 ± 0.063		
Cu II	1358.8	< 12.793	< 12.793	< -0.847
Zn II	2026.1	13.036 ± 0.022	13.035 ± 0.022	-0.985 ± 0.102
Zn II	2062.7	< 13.100		

4.8. FJ0812+32, $z = 2.626$

This FIRST quasar shows the remarkable damped Ly α system at $z = 2.626$ which exhibits among the strongest metal-line transitions of any damped Ly α system. In fact, our follow-up HIRES observations revealed the first discovery of Ge⁺, Cl⁰, B⁺, and several other ions (Prochaska, Howk, & Wolfe 2003). The Ly α profile from our ESI spectrum is presented in Figure 21. The fit is somewhat challenging because of the proximity to the Ly α peak and the gap in data quality at $\lambda \approx 4475\text{\AA}$ owing to a defect on the CCD. Nevertheless, the core is well sampled and the blue wing provides complimentary constraints.

An extensive list of metal-line transitions are plotted in Figure 22 and the corresponding ionic column densities are provided in Table 14. A comprehensive analysis of this sightline including this ESI data will be presented in a future paper.

4.9. Q0930+28, $z = 3.235$




FIG. 21.— Ly α profile of the damped Ly α system at $z = 2.626$ toward FJ0812+32. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.35^{+0.10}_{-0.10}$. This Ly α profile is complicated by the chip defect at $\lambda \approx 4470\text{\AA}$ and the presence of the Ly α emission peak. The core, however, is well sampled and the fit is precise.

TABLE 15
IONIC COLUMN DENSITIES: Q0930+28, $z = 3.235$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1036.3	> 14.173	> 14.173	> -2.767
C II	1334.5	> 13.887		
C II*	1335.7	< 12.854		
C IV	1548.2	13.210 ± 0.059		
C IV	1550.8	< 13.152		
O I	1302.2	> 14.196	> 14.196	> -2.894
Al II	1670.8	12.369 ± 0.026	12.369 ± 0.026	-2.471 ± 0.103
Si II	1304.4	13.793 ± 0.022	13.793 ± 0.022	-2.117 ± 0.102
Si II	1526.7	> 13.713		
Si IV	1393.8	12.749 ± 0.053		
Si IV	1402.8	12.749 ± 0.101		
Fe II	1144.9	13.389 ± 0.079	13.490 ± 0.031	-2.360 ± 0.105
Fe II	1608.5	13.540 ± 0.040		
Fe II	2344.2	13.479 ± 0.062		
Fe II	2374.5	< 13.713		

This damped Ly α system and its neighboring sub-DLA ($z = 3.246$) have been studied previously in the literature (Lu et al. 1998; Prochaska et al. 2002). Our observations provide greater wavelength coverage and are generally consistent with the echelle observations. The Ly α profiles of the DLA and sub-DLA are shown in Figure 23 while the metal-line profiles and ionic column densities for the DLA are given by Figure 24 and Table 15.

4.10. PC0953+47, $z = 3.404, 3.891, 4.244$

This quasar discovered by Schneider et al. (1991) and one could identify three $z > 3$ DLA candidates in their discovery spectrum. The systems are particularly unusual because all three DLA have among the largest HI column densities at these redshifts. The Ly α profiles are shown individually in Figures 25-27. The S/N of this spectrum is among the lowest in our dataset, primarily because of




FIG. 22.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.626$ toward FJ0812+32. The vertical line at $v = 0$ corresponds to $z = 2.62644$.




Fig 22 – cont




Fig 22 – cont




FIG. 23.— Ly α profile of the damped Ly α system at $z = 3.235$ toward Q0930+28. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.35^{+0.10}_{-0.10}$. This solution is complicated by the presence of a neighboring sub-DLA ($z = 3.246$, $N(\text{HI}) \approx 10^{20.2} \text{ cm}^{-2}$) although we consider the final value to be relatively accurate.

TABLE 16

IONIC COLUMN DENSITIES: PC0953+47, $z = 3.404$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1550.8	> 14.858		
Al II	1670.8	> 13.360	> 13.360	> -2.280
Si II	1526.7	> 14.617	> 14.617	> -2.093
Si II	1808.0	< 15.166		
Fe II	1608.5	> 14.472	> 14.472	> -2.178
Fe II	1611.2	< 15.303		
Ni II	1709.6	< 13.896	< 13.896	< -1.504

TABLE 17

IONIC COLUMN DENSITIES: PC0953+47, $z = 3.891$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	> 14.755		
C IV	1550.8	> 14.687		
Al II	1670.8	> 13.831	> 13.831	> -1.859
Si II	1526.7	> 15.161	> 15.161	> -1.599
Si II	1808.0	< 15.364		
Si IV	1393.8	> 14.138		
Si IV	1402.8	> 13.988		
Fe II	1608.5	> 14.988	> 14.988	> -1.712
Fe II	1611.2	< 15.530		
Ni II	1370.1	13.732 ± 0.062	13.752 ± 0.055	-1.698 ± 0.114
Ni II	1741.5	13.859 ± 0.115		
Ni II	1751.9	< 14.055		

poor seeing during this particular observation. For the $z = 3.404$ and $z = 4.244$ DLA, the continuum is poorly constrained and we place a larger uncertainty on their H I column densities.

The metal-line profiles for the three DLA are presented in Figures 28-30 and the ionic column densities are listed in Tables 16-18. Unfortunately, the DLA at $z = 3.404$ ex-




FIG. 24.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.235$ toward Q0930+28. The vertical line at $v = 0$ corresponds to $z = 3.2352$.

TABLE 18

IONIC COLUMN DENSITIES: PC0953+47, $z = 4.244$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 14.719	> 14.719	> -2.771
C II*	1335.7	13.599 ± 0.099		
C IV	1548.2	14.273 ± 0.052		
C IV	1550.8	14.217 ± 0.092		
O I	1302.2	> 15.162	> 15.162	> -2.478
Al II	1670.8	< 12.873	< 12.873	< -2.517
Si II	1304.4	14.235 ± 0.027	14.235 ± 0.027	-2.225 ± 0.152
Si II	1526.7	> 14.104		
Fe II	1608.5	13.900 ± 0.068	13.900 ± 0.068	-2.500 ± 0.165
Fe II	1611.2	< 15.080		
Ni II	1370.1	< 13.613	< 13.613	< -1.537
Ni II	1741.5	< 13.821		
Ni II	1751.9	< 14.050		




FIG. 25.— Ly α profile of the damped Ly α system at $z = 3.404$ toward PC0953+47. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.15^{+0.15}_{-0.15}$. The continuum (dotted line) is not well constrained in this region of the spectra and the signal-to-noise ratio is poor. Therefore, we ascribe a higher uncertainty to the $N(\text{HI})$ value of this DLA. Also note that we were forced to include an additional absorber at $\lambda \approx 5390\text{\AA}$ to explain the complete core of this Ly α profile.




FIG. 27.— Ly α profile of the damped Ly α system at $z = 4.244$ toward PC0953+47. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.90^{+0.15}_{-0.15}$. There is significant contamination from the Ly α forest in the region surrounding this DLA and the continuum is complicated at the red end by the Ly α emission peak. The resulting fit to the Ly α profile is quite uncertain. Note the feature at $6362\text{\AA}$ is a skyline as denoted by the spike in the sigma spectrum.




FIG. 26.— Ly α profile of the damped Ly α system at $z = 3.891$ toward PC0953+47. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.20^{+0.10}_{-0.10}$. In comparison with the DLA at $z = 3.404$, the continuum is better determined across this DLA and the resulting solution is nicely constrained.

hibits transitions which are either too weak or too strong for an accurate measure of the ionic column density for the few transitions which lie outside the Ly α forest. This is nearly the case for the DLA at $z = 3.891$ as well. Because of the remarkable nature of this sightline, we hope to achieve higher S/N with future observations.

TABLE 19
IONIC COLUMN DENSITIES: PSS0957+33, $z = 3.280$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	13.945 ± 0.015		
C IV	1550.8	13.838 ± 0.033		
N I	1200.7	< 14.444	< 14.444	< -1.936
Al II	1670.8	> 13.128	> 13.128	> -1.812
Al III	1854.7	12.544 ± 0.042		
Al III	1862.8	12.396 ± 0.099		
Si II	1526.7	> 14.461	14.676 ± 0.078	-1.334 ± 0.127
Si II	1808.0	14.676 ± 0.078		
S II	1259.5	< 14.577	< 14.577	< -1.073
Cr II	2056.3	< 12.891	< 12.891	< -1.229
Fe II	1608.5	< 14.261	> 14.088	> -1.862
Fe II	2249.9	< 14.821		
Fe II	2260.8	< 14.575		
Fe II	2344.2	> 14.088		
Fe II	2374.5	< 14.102		
Ni II	1741.5	< 13.152	< 13.152	< -1.548
Zn II	2026.1	< 12.168	< 12.168	< -0.952

4.11. PSS0957+33, $z = 3.280, 4.180$

This sightline was first analysed in PGW01 and is notable for exhibiting two DLA at $z > 3$. Our re-analysis has revised the HI values of the DLA upward by ≈ 0.1 dex primarily due to a reassessment of the continuum (Figures 31,32). The velocity plots for the metal-line transitions are given by Figures 33 and 34 and the ionic column densities are listed in Tables 19 and 20. We note in passing that our comparison of the ESI data with our HIRES observations (Prochaska et al. 2001) suggest significant saturation in most of the ESI profiles of the $z = 4.18$ DLA.

4.12. BQ1021+3001, $z = 2.949$




FIG. 28.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.404$ toward PC0953+47. The vertical line at $v = 0$ corresponds to $z = 3.4036$.

TABLE 20

IONIC COLUMN DENSITIES: PSS0957+33, $z = 4.180$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 14.799	> 14.799	> -2.441
C IV	1548.2	< 14.081		
C IV	1550.8	13.910 ± 0.031		
O I	1302.2	> 14.972	> 14.972	> -2.418
Al II	1670.8	13.038 ± 0.015	13.038 ± 0.015	-2.102 ± 0.151
Al III	1854.7	12.775 ± 0.103		
Al III	1862.8	< 12.878		
Si II	1304.4	> 14.376	> 14.376	> -1.834
Si II	1526.7	> 14.250		
Si IV	1393.8	12.931 ± 0.052		
Si IV	1402.8	13.401 ± 0.035		
Fe II	1608.5	> 13.897	> 13.897	> -2.253
Fe II	1611.2	< 15.038		
Ni II	1317.2	< 12.948	< 12.948	< -1.952

This quasar was drawn from the FIRST bright quasar survey (Gregg et al. 1996) and White et al. (2000) identified this DLA as a candidate in their follow-up observations. Our fit to the Ly α profile is shown in Figure 35, Figure 36 presents the velocity profiles, and the ionic column densities are listed in Table 21.




FIG. 29.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.891$ toward PC0953+47. The vertical line at $v = 0$ corresponds to $z = 3.8910$.

4.13. CTQ460, $z = 2.777$

This DLA was discovered by Lopez et al. (2001) toward the quasar CTQ460 which is drawn from the Calán-Tololo survey (Maza et al. 1995). The Ly α profile presented in Figure 37 has a well sampled core and a relatively precise H I measurement. Figure 38 provides an exhaustive list of metal-line profiles and Table 22 gives the ionic column densities.




FIG. 30.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 4.244$ toward PC0953+47. The vertical line at $v = 0$ corresponds to $z = 4.24419$.




FIG. 31.— Ly α profile of the damped Ly α system at $z = 3.280$ toward PSS0957+33. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.45^{+0.10}_{-0.10}$. Aside from the somewhat significant amount of contamination from coincident Ly α clouds, this profile offered a straightforward solution.




FIG. 32.— Ly α profile of the damped Ly α system at $z = 4.180$ toward PSS0957+33. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.65^{+0.15}_{-0.15}$. Both the presence of the nearby Ly α emission peak and the absence of detailed information in the core of the profile lead to a more uncertain HI solution.




FIG. 33.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.280$ toward PSS0957+33. The vertical line at $v = 0$ corresponds to $z = 3.2797$.




FIG. 34.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 4.180$ toward PSS0957+33. The vertical line at $v = 0$ corresponds to $z = 4.179825$.

TABLE 21

IONIC COLUMN DENSITIES: BQ1021+3001, $z = 2.949$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.9	< 12.772		
C II	1334.5	> 14.586	> 14.586	> -2.704
C II*	1335.7	< 12.909		
C IV	1548.2	13.991 ± 0.013		
C IV	1550.8	13.957 ± 0.025		
O I	1302.2	> 14.848	> 14.848	> -2.592
Al II	1670.8	12.693 ± 0.022	12.693 ± 0.022	-2.497 ± 0.102
Al III	1854.7	< 12.377		
Si II	1304.4	14.090 ± 0.017	14.090 ± 0.017	-2.170 ± 0.101
Si II	1526.7	> 14.003		
Si II	1808.0	< 14.603		
Si IV	1393.8	> 13.602		
Si IV	1402.8	13.585 ± 0.029		
Cr II	2056.3	< 12.833	< 12.833	< -1.537
Fe II	1608.5	13.880 ± 0.026	13.880 ± 0.026	-2.320 ± 0.103
Fe II	2344.2	> 13.636		
Fe II	2382.8	> 13.561		
Ni II	1370.1	< 13.368	< 13.368	< -1.582
Ni II	1709.6	< 13.400		
Zn II	2026.1	< 12.189	< 12.189	< -1.181




FIG. 35.— Ly α profile of the damped Ly α system at $z = 2.949$ toward BQ1021+3001. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.70^{+0.10}_{-0.10}$. Although continuum placement (dotted line) was particularly challenging across this portion of the spectrum, the core is rather free from line blending and the resulting fit is well constrained.




FIG. 36.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.949$ toward BQ1021+3001. The vertical line at $v = 0$ corresponds to $z = 2.9489$.




Fig 36 – cont

TABLE 22
IONIC COLUMN DENSITIES: CTQ460, $z = 2.777$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 15.151	> 15.151	> -2.439
C II*	1335.7	12.969 ± 0.051		
C IV	1548.2	14.016 ± 0.011		
C IV	1550.8	14.051 ± 0.017		
O I	1302.2	> 15.564	> 15.564	> -2.176
Mg II	1239.9	< 15.054	< 15.054	< -1.526
Mg II	1240.4	< 15.292		
Al II	1670.8	13.425 ± 0.007	13.425 ± 0.007	-2.065 ± 0.100
Al III	1854.7	13.025 ± 0.017		
Al III	1862.8	13.057 ± 0.029		
Si II	1304.4	> 14.970	> 14.970	> -1.590
Si II	1526.7	> 14.726		
Si IV	1393.8	13.774 ± 0.008		
Si IV	1402.8	13.852 ± 0.013		
S II	1259.5	14.790 ± 0.022	14.791 ± 0.022	-1.409 ± 0.102
Cr II	2056.3	13.008 ± 0.054	13.008 ± 0.054	-1.662 ± 0.114
Mn II	2576.9	12.662 ± 0.054	12.662 ± 0.054	-1.868 ± 0.114
Fe II	1608.5	14.683 ± 0.008	14.684 ± 0.005	-1.816 ± 0.100
Fe II	1611.2	< 14.612		
Fe II	2249.9	< 14.758		
Fe II	2260.8	14.657 ± 0.056		
Fe II	2344.2	> 14.477		
Fe II	2374.5	14.686 ± 0.008		
Fe II	2382.8	> 14.258		
Fe II	2586.7	> 14.503		
Fe II	2600.2	> 14.181		
Ni II	1370.1	13.572 ± 0.032	13.587 ± 0.024	-1.663 ± 0.103
Ni II	1454.8	13.447 ± 0.115		
Ni II	1709.6	13.573 ± 0.064		
Ni II	1741.6	13.675 ± 0.060		
Ni II	1751.9	13.676 ± 0.061		
Zn II	2062.7	< 12.359	< 12.359	< -1.311




FIG. 37.— Ly α profile of the damped Ly α system at $z = 2.777$ toward CTQ460. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.00^{+0.10}_{-0.10}$. Although the continuum is not particularly well constrained across this DLA and the blue wing approaches the CCD defect, the core is well sampled and constrains the HI column density.




FIG. 38.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.777$ toward CTQ460. The vertical line at $v = 0$ corresponds to $z = 2.7775$.

Fig 38 – cont

TABLE 23

IONIC COLUMN DENSITIES: HS1132+2243, $z = 2.783$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.9	< 12.569		
C II	1334.5	> 14.580	> 14.580	> -3.010
C II*	1335.7	< 12.931		
C IV	1548.2	13.370 ± 0.036		
C IV	1550.8	13.365 ± 0.070		
N I	1199.5	14.127 ± 0.020	14.019 ± 0.019	-2.911 ± 0.073
N I	1200.2	13.863 ± 0.039		
N II	1084.0	< 13.431		
O I	1302.2	> 14.989	> 14.989	> -2.751
Al II	1670.8	> 12.867	> 12.867	> -2.623
Al III	1854.7	12.141 ± 0.107		
Si II	1193.3	> 14.015	14.491 ± 0.118	-2.069 ± 0.137
Si II	1260.4	> 13.890		
Si II	1304.4	> 14.303		
Si II	1526.7	> 14.186		
Si II	1808.0	14.491 ± 0.118		
Si IV	1393.8	13.008 ± 0.039		
Si IV	1402.8	13.000 ± 0.081		
S II	1250.6	< 14.113	14.071 ± 0.060	-2.129 ± 0.092
S II	1259.5	14.071 ± 0.060		
Cr II	2056.3	12.834 ± 0.104	12.834 ± 0.104	-1.836 ± 0.125
Mn II	2576.9	< 12.457	< 12.457	< -2.073
Fe II	1144.9	13.951 ± 0.044	14.024 ± 0.014	-2.476 ± 0.071
Fe II	1608.5	14.084 ± 0.019		
Fe II	1611.2	< 14.504		
Fe II	2260.8	< 14.386		
Fe II	2344.2	> 13.870		
Fe II	2374.5	14.082 ± 0.027		
Fe II	2382.8	> 13.652		
Fe II	2586.7	13.900 ± 0.035		
Fe II	2600.2	> 13.596		
Fe III	1122.5	< 13.726		
Ni II	1317.2	< 13.155	< 13.155	< -2.095
Ni II	1370.1	< 13.164		
Ni II	1741.5	< 13.249		
Ni II	1751.9	< 13.402		
Zn II	2026.1	< 11.986	< 11.986	< -1.684

4.14. *HS1132+2243*, $z = 2.783$

This quasar is a member of the Hamburg-ESO Quasar Survey (Hagen et al. 1999) and one could identify the DLA in their discovery spectrum. Although the red wing of the Ly α profile occurs along the Ly α emission peak of the quasar, the $N(\text{HI})$ is very well set by the core of the Ly α profile (Figure 39). The extensive set of metal-line transitions are given in Figure 40 and Table 23 lists the ionic column densities.

4.15. *Q1209+0919*, $z = 2.584$

This DLA is a member of the LBQS survey of Wolfe et al. (1995) and exhibits a relatively large HI column density. It has one of the lowest redshifts in this survey and was chosen primarily because of its large HI column density. Our fit to the Ly α profile is presented in Figure 41. Figure 42 plots the velocity profiles which include Zn⁺ and Mn⁺ detections. Finally, the ionic column densities are provided in Table 24.

4.16. *PSS1248+31*, $z = 3.696$

This PSS quasar was first analysed in PGW01 and we note that our revised fit to the Ly α profile reveals a significantly larger HI column density (Figure 43) owing primarily to a re-evaluation of the quasar continuum. In PGW01,




FIG. 39.— Ly α profile of the damped Ly α system at $z = 2.783$ toward HS1132+2243. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.00^{+0.07}_{-0.07}$. Although the red wing of the Ly α profile occurs along the Ly α emission peak of the quasar, the $N(\text{HI})$ is very well determined by the core of the Ly α profile.

TABLE 24

IONIC COLUMN DENSITIES: Q1209+0919, $z = 2.584$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1560.3	< 13.192		
C I	1656.9	13.276 ± 0.085		
C IV	1548.2	> 14.976		
C IV	1550.8	> 14.973		
Al II	1670.8	> 14.068	> 14.068	> -1.822
Al III	1854.7	13.712 ± 0.013		
Al III	1862.8	13.826 ± 0.017		
Si II	1526.7	> 15.295	> 15.990	> -0.970
Si II	1808.0	> 15.990		
Ti II	1910.6	< 12.652	< 12.652	< -1.688
Cr II	2056.3	13.609 ± 0.053	13.581 ± 0.050	-1.489 ± 0.112
Cr II	2066.2	13.468 ± 0.137		
Mn II	2576.9	13.134 ± 0.059	13.142 ± 0.049	-1.788 ± 0.111
Mn II	2594.5	13.159 ± 0.084		
Fe II	1608.5	> 15.004	15.219 ± 0.044	-1.681 ± 0.109
Fe II	1611.2	< 15.043		
Fe II	2249.9	< 15.283		
Fe II	2260.8	15.219 ± 0.044		
Fe II	2344.2	> 14.894		
Fe II	2374.5	> 15.324		
Fe II	2382.8	> 14.730		
Ni II	1703.4	< 14.505	< 14.218	< -1.432
Ni II	1751.9	< 14.218		
Zn II	2026.1	12.982 ± 0.053	12.982 ± 0.053	-1.088 ± 0.113

we fit the continuum prior to fitting the Ly α profile and in the process significantly underestimated both the continuum and the $N(\text{HI})$ value. We consider the new solution to be very well constrained by the observations. Figure 44 gives the metal-line profiles and Table 25 lists the ionic column densities.

4.17. *PSS1253-0228*, $z = 2.783$

This quasar sightline is remarkable because it may exhibit the largest $N(\text{HI})$ value of any known high z damped




FIG. 40.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.783$ toward HS1132+2243. The vertical line at $v = 0$ corresponds to $z = 2.78347$.




Fig 40 – cont

system. Figure 45 presents the Ly α profile and our best fit to the data assuming $N(\text{HI}) = 10^{21.85} \text{ cm}^{-2}$. The fit is complicated by several factors: (1) the coincidence with the CCD defect of ESI at $\lambda \approx 4500 \text{ \AA}$; (2) somewhat poor S/N; (3) a rather uncertain continuum; and (4) blending with a sub-DLA at $z = 3.603$ which is included in our analysis. Nevertheless, it is difficult to fit the core of this profile with an HI value significantly less than what we have adopted. Granted the special nature of this DLA, future observations using an instrument with higher efficiency at $\lambda \approx 4400 - 4800 \text{ \AA}$ are warranted. We stress that Peroux et al. (2001) reported a significantly lower value to $N(\text{HI})$ for this DLA.

Figure 46 presents the few metal-line transitions outside the Ly α forest. Note that SiII 1808 is clearly saturated while Zn II 2026 is blended with coincident absorption features. We do estimate the Zn^+ column density, however,

TABLE 25
IONIC COLUMN DENSITIES: PSS1248+31, $z = 3.696$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	13.841 ± 0.024		
C IV	1550.8	13.903 ± 0.047		
Al II	1670.8	12.777 ± 0.020	12.777 ± 0.020	-2.343 ± 0.073
Al III	1854.7	< 12.191		
Si II	1304.4	14.393 ± 0.016	14.393 ± 0.016	-1.797 ± 0.072
Si II	1526.7	> 14.243		
Si II	1808.0	< 14.610		
Si IV	1393.8	13.414 ± 0.017		
Si IV	1402.8	13.310 ± 0.045		
Cr II	2056.3	< 13.242	< 13.242	< -1.058
Fe II	1608.5	13.890 ± 0.046	13.890 ± 0.046	-2.240 ± 0.084
Ni II	1454.8	< 13.622	< 13.493	< -1.387
Ni II	1741.6	< 13.493		
Zn II	2026.1	< 12.615	< 12.615	< -0.685




FIG. 41.— Ly α profile of the damped Ly α system at $z = 2.584$ toward Q1209+0919. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.40^{+0.10}_{-0.10}$. The fit is complicated by the poor behavior of the continuum across the DLA which includes the QSO OVI emission feature at $\lambda \approx 4450\text{\AA}$, but the value is relatively well constrained by the core of the Ly α profile.

TABLE 26

IONIC COLUMN DENSITIES: PSS1253-0228, $z = 2.783$

Ion	λ	AODM	N_{adopt}	$[X/H]$
C II	1334.5	> 15.012	> 15.012	> -3.428
O I	1302.2	> 15.347	> 15.347	> -3.243
Al II	1670.8	> 13.399	> 13.399	> -2.941
Al III	1854.7	12.537 ± 0.097		
Al III	1862.8	< 12.664		
Si II	1808.0	> 15.597	> 15.597	> -1.813
Si IV	1402.8	13.672 ± 0.052		
Ti II	1910.6	< 12.841	< 12.842	< -1.948
Cr II	2056.3	13.693 ± 0.029	13.632 ± 0.025	-1.888 ± 0.202
Cr II	2062.2	13.537 ± 0.048		
Mn II	2576.9	13.420 ± 0.070	13.420 ± 0.070	-1.960 ± 0.212
Fe II	1608.5	> 14.627	15.358 ± 0.038	-1.992 ± 0.204
Fe II	1611.2	15.340 ± 0.049		
Fe II	2249.9	15.393 ± 0.061		
Fe II	2260.8	> 15.207		
Fe II	2344.2	> 14.331		
Fe II	2374.5	> 14.752		
Fe II	2382.8	> 14.136		
Fe II	2586.7	> 14.388		
Fe II	2600.2	> 14.014		
Ni II	1709.6	14.359 ± 0.048	14.170 ± 0.036	-1.930 ± 0.203
Ni II	1741.5	14.050 ± 0.069		
Ni II	1751.9	14.149 ± 0.072		
Zn II	2026.1	< 13.068	12.768 ± 0.074	-1.752 ± 0.213
Zn II	2062.7	12.768 ± 0.074		

from the unblended portion of the Zn II 2062 profile (apparent as the dotted line in the Cr II 2062 profile). Table 26 provides the column density measurements for this DLA.




FIG. 42.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.584$ toward Q1209+0919. The vertical line at $v = 0$ corresponds to $z = 2.5841$.




Fig 42 – cont




FIG. 43.— Ly α profile of the damped Ly α system at $z = 3.696$ toward PSS1248+31. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.63^{+0.07}_{-0.07}$. Because of minimal contamination within the core of the Ly α profile and a well behaved continuum, we are particularly certain of the HI value for this DLA.




FIG. 44.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.696$ toward PSS1248+31. The vertical line at $v = 0$ corresponds to $z = 3.6960$.




FIG. 45.— Ly α profile of the damped Ly α system at $z = 2.783$ toward PSS1253-0228. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.85^{+0.20}_{-0.20}$. Although the fit is complicated by several factors (see the text), we consider this value to be relatively secure. In particular, it is difficult for the HI column density to be significantly lower than $10^{21.85} \text{ cm}^{-2}$.




FIG. 46.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.783$ toward PSS1253-0228. The vertical line at $v = 0$ corresponds to $z = 2.78282$.

Fig 46 – cont

TABLE 27

IONIC COLUMN DENSITIES: Q1337+11, $z = 2.795$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 14.483	> 14.483	> -3.057
C II*	1335.7	< 13.093		
C IV	1548.2	13.476 ± 0.050		
C IV	1550.8	13.508 ± 0.089		
N I	1199.5	13.751 ± 0.060	13.751 ± 0.060	-3.129 ± 0.117
N II	1084.0	< 13.550		
O I	1302.2	> 14.866	> 14.866	> -2.824
Al II	1670.8	12.844 ± 0.021	12.844 ± 0.021	-2.596 ± 0.102
Al III	1854.7	12.546 ± 0.069		
Al III	1862.8	< 12.513		
Si II	1260.4	> 13.568	14.718 ± 0.116	-1.792 ± 0.153
Si II	1304.4	> 14.274		
Si II	1526.7	> 14.202		
Si II	1808.0	14.719 ± 0.116		
Si III	1206.5	> 13.488		
Si IV	1393.8	12.894 ± 0.065		
Si IV	1402.8	< 12.861		
S II	1259.5	14.270 ± 0.035	14.270 ± 0.036	-1.880 ± 0.106
Cr II	2056.3	< 12.900	< 12.900	< -1.720
Mn II	2576.9	< 12.528	< 12.528	< -1.952
Fe II	1608.5	14.051 ± 0.021	14.062 ± 0.018	-2.388 ± 0.102
Fe II	1611.2	< 14.767		
Fe II	2260.8	< 14.487		
Fe II	2344.2	> 13.897		
Fe II	2374.5	14.101 ± 0.037		
Fe II	2382.8	> 13.634		
Fe II	2586.7	13.979 ± 0.052		
Fe II	2600.2	> 13.575		
Ni II	1370.1	< 13.199	< 13.199	< -2.001
Ni II	1741.6	< 13.424		
Zn II	2026.1	< 12.201	< 12.201	< -1.419




FIG. 47.— Ly α profile of the damped Ly α system at $z = 2.795$ toward Q1337+11. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.95^{+0.10}_{-0.10}$. The red wing continuum is complicated by the Ly α emission peak, but the fit is secure.

4.18. Q1337+11, $z = 2.795$

This LBQS quasar (Wolfe et al. 1995) was discovered in the original surveys of Sargent, Steidel & Boksenberg (1989) and was chosen principally to fill the gap at $z \approx 2.8$




FIG. 48.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.795$ toward Q1337+11. The vertical line at $v = 0$ corresponds to $z = 2.79545$.

of previous metallicity surveys. The Ly α profile of this DLA is shown in Figure 47 along with the best fit solution. Although the Ly α emission peak complicates the solution, the fit is reasonably secure. The metal-line profiles are shown in Figure 48 and the ionic column densities are listed in Table 27. Note the Zn II 2026 profile is blended with atmospheric absorption and can only be considered an upper limit to $N(\text{Zn}^+)$.

4.19. PKS1354-17, $z = 2.780$

This radio-bright quasar was first discovered by Schilizzi & McAdam (1975) and we discovered the damped Ly α as part of back-up observing plan on the Magellan telescope. Figure 49 plots the Ly α profile and solution for this damped system. Because of the very poor S/N, we report an uncertainty of 0.15 dex and caution that this system may not meet the DLA criterion of $2 \times 10^{20} \text{ cm}^{-2}$ when




Fig 48 – cont

TABLE 28

IONIC COLUMN DENSITIES: PKS1354-17, $z = 2.780$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 14.142	> 14.142	> -2.748
C II*	1335.7	< 12.833		
C IV	1548.2	13.670 ± 0.123		
C IV	1550.8	< 13.857		
Al II	1670.8	12.395 ± 0.095	12.395 ± 0.095	-2.395 ± 0.178
Si II	1526.7	13.976 ± 0.056	13.976 ± 0.056	-1.884 ± 0.160
Si II	1808.0	< 14.870		
Si IV	1402.8	< 13.347		
Fe II	1608.5	< 13.881	13.374 ± 0.080	-2.426 ± 0.170
Fe II	2344.2	13.374 ± 0.080		
Fe II	2374.5	< 13.761		

additional observations are made. Figure 50 plots the few metal-line transitions observed and Table 28 provides the measurements of ionic column densities.




FIG. 49.— Ly α profile of the damped Ly α system at $z = 2.780$ toward PKS1354-17. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.30^{+0.15}_{-0.15}$. The signal-to-noise ratio is very poor in this portion of the spectrum and we report an uncertainty of 0.15 dex in the fit.

TABLE 29

IONIC COLUMN DENSITIES: PSS1432+39, $z = 3.272$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.9	< 13.047		
C IV	1548.2	13.751 ± 0.030		
C IV	1550.8	13.637 ± 0.075		
Al II	1670.8	> 13.749	> 13.749	> -1.991
Al III	1854.7	12.920 ± 0.098		
Si II	1526.7	> 15.062	15.666 ± 0.048	-1.144 ± 0.111
Si II	1808.0	15.666 ± 0.048		
Cr II	2056.3	< 13.160	< 13.160	< -1.760
Fe II	1608.5	> 14.930	> 14.930	> -1.820
Fe II	1611.2	< 15.070		
Fe II	2249.9	< 15.312		
Ni II	1709.6	< 13.779	13.750 ± 0.101	-1.750 ± 0.142
Ni II	1741.6	13.750 ± 0.101		
Ni II	1751.9	< 13.834		
Zn II	2026.1	< 12.645	< 12.645	< -1.275

4.20. *PSS1432+39*, $z = 3.272$

We originally discovered and analysed this damped Ly α system in PGW01. Similar to other DLA from the PGW01 sample, we find a somewhat higher HI value in this analysis (Figure 51). Again, this is due to the improved approach we have taken to measuring HI column densities in the present work. The metal-line profiles for this DLA are given in Figure 52 and the ionic column densities are listed in Table 29. For this DLA, the Ni II 1709, 1741, and 1751 profiles are all marginally detected and yield a Ni⁺ column density of $10^{13.75} \text{ cm}^{-2}$ when treated together.

4.21. *Q1502+4837*, $z = 2.570$

This quasar is a member of the Cosmic Lens All Sky Survey (CLASS) (Snellen et al. 2001) and the DLA at $z = 2.57$ was identified in their discovery spectrum. The fit to the Ly α profile (Figure 53) is complicated by the




FIG. 50.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.780$ toward PKS1354-17. The vertical line at $v = 0$ corresponds to $z = 2.77993$.

TABLE 30

IONIC COLUMN DENSITIES: Q1502+4837, $z = 2.570$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 15.174	> 15.174	> -1.716
C IV	1548.2	14.240 ± 0.034		
C IV	1550.8	14.149 ± 0.067		
O I	1302.2	> 15.341	> 15.341	> -1.699
Mg II	2796.4	> 13.961	> 13.961	> -1.919
Al II	1670.8	> 13.361	> 13.362	> -1.428
Al III	1854.7	12.887 ± 0.144		
Si II	1526.7	14.241 ± 0.085	14.241 ± 0.085	-1.619 ± 0.172
Si IV	1393.8	> 14.434		
Fe II	1608.5	14.151 ± 0.121	14.151 ± 0.121	-1.649 ± 0.193
Fe II	2344.2	14.103 ± 0.064		
Fe II	2374.5	< 14.011		
Fe II	2600.2	> 14.081		




FIG. 51.— Ly α profile of the damped Ly α system at $z = 3.272$ toward PSS1432+39. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.25^{+0.10}_{-0.10}$. Note this value is a factor of 0.3 dex higher than the value listed in PGW01. This is primarily due to our decision to fit fluxed data, i.e. including the continuum in the solution when fitting the profile as described in § 3.

poor efficiency of ESI at $\lambda < 4400\text{\AA}$, but the fit is reasonable. We note, however, that future observations may favor a central value below the DLA criterion. Figure 54 provides the metal-line transitions and Table 30 gives the ionic column densities. The S/N of this spectrum is poor and only a few profiles provide meaningful measurements. We caution using this DLA in ones analysis. Finally, we note in passing that the velocity profiles span an impressive 400 km/s.

TABLE 31

IONIC COLUMN DENSITIES: PSS1506+5220, $z = 3.224$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.9	< 12.787		
C II	1334.5	> 15.210	> 15.210	> -2.050
C II*	1335.7	< 12.890		
C IV	1548.2	13.581 ± 0.033		
C IV	1550.8	13.746 ± 0.048		
O I	1302.2	> 14.656	> 14.657	> -2.753
Al II	1670.8	> 12.504	> 12.503	> -2.657
Al III	1854.7	< 12.080		
Si II	1260.4	> 13.536	13.882 ± 0.018	-2.348 ± 0.072
Si II	1526.7	13.882 ± 0.018		
Si III	1206.5	> 13.379		
Cr II	2056.3	< 12.784	< 12.784	< -1.556
Fe II	1608.5	13.896 ± 0.036	13.708 ± 0.034	-2.462 ± 0.078
Fe II	2344.2	13.546 ± 0.062		
Ni II	1709.6	< 13.413	< 13.413	< -1.507
Zn II	2026.1	< 12.105	< 12.105	< -1.235

4.22. PSS1506+5220, $z = 3.224$

This PSS quasar exhibits a damped system at $z = 3.224$ with a reasonably simple Ly α profile (Figure 55). Because




FIG. 52.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.272$ toward PSS1432+39. The vertical line at $v = 0$ corresponds to $z = 3.2720$.

the continuum is also well behaved, we place an uncertainty on the $N(\text{HI})$ value of only 0.07 dex. Figure 56 plots the velocity profiles of this low metallicity, simple DLA and the ionic column densities are given in Table 31.

4.23. PSS1723+2243, $z = 3.695$

The damped Ly α system toward this Palomar Sky Survey quasar is remarkable for exhibiting a particularly large metallicity at $z > 3.5$ as well as metal-line profiles which span over 400 km/s. The HI column density is best fit with two components separated by ≈ 300 km s $^{-1}$ as shown in Figure 57. The $N(\text{HI})$ values for the individual components is not very well constrained and if one divides the total HI column density evenly, then neither absorber satisfies the DLA criterion of $10^{20.3}$ cm $^{-2}$. In this case, one might refer to this damped system as a pair of sub-DLA. To further complicate matters, if the Ly α profile




FIG. 53.— Ly α profile of the damped Ly α system at $z = 2.570$ toward Q1502+4837. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.30^{+0.15}_{-0.15}$. Although the profile is relatively clean, we report an uncertainty of 0.15 dex because of the poor S/N.

TABLE 32
IONIC COLUMN DENSITIES: PSS1723+2243, $z = 3.695$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	> 14.475		
C IV	1550.8	14.761 ± 0.007		
Al II	1670.8	> 13.564	> 13.564	> -1.426
Al III	1862.8	< 12.603		
Si II	1526.7	> 14.861	> 14.861	> -1.199
Si II	1808.0	< 15.497		
Cr II	2056.3	< 13.214	< 13.214	< -0.956
Fe II	1608.5	> 14.575	> 14.575	> -1.425
Ni II	1741.6	< 13.950	< 13.950	< -0.800
Zn II	2026.1	> 12.508	> 12.508	> -0.662

were observed at lower resolution, one may have been likely compose a fit with a single component with $N(\text{HI}) \approx 10^{20.5}$ cm $^{-2}$. Unfortunately, a higher redshift Lyman limit system precludes an analysis of the higher order Lyman series for the system at $z = 3.695$. For now, we fit the profile with two components and report the total $N(\text{HI})$ value. Similarly, we have chosen to integrate the metal-line profiles over the full velocity range. Any future studies which focus on the relative abundances of this system must take caution in interpreting the metallicity of the gas.

The metal-line profiles for this DLA are shown in Figure 58 which includes only a few species because of bad line blending both within and outside the Ly α forest. The ionic column densities are provided in Table 32. We have cautiously listed most of the values as lower limits because our experience with ESI suggests several of the line profiles are saturated.




FIG. 54.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.570$ toward Q1502+4837. The vertical line at $v = 0$ corresponds to $z = 2.56956$.




FIG. 55.— Ly α profile of the damped Ly α system at $z = 3.224$ toward PSS1506+5220. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.67^{+0.07}_{-0.07}$. The Ly α profile has a poorly determined blue wing but sufficient sampling in its core and red wing to provide a rather precise fit.

TABLE 33

IONIC COLUMN DENSITIES: PSS2155+1358, $z = 3.316$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	> 14.221		
C IV	1550.8	14.209 ± 0.018		
Al II	1670.8	> 13.316	> 13.316	> -1.724
Si II	1526.7	> 14.649	14.847 ± 0.072	-1.263 ± 0.166
Si II	1808.0	14.848 ± 0.072		
Cr II	2056.3	< 13.150	< 13.151	< -1.069
Fe II	1608.5	> 14.373	> 14.373	> -1.677
Fe II	2249.9	< 15.124		
Fe II	2344.2	> 14.284		
Ni II	1709.6	< 13.437	< 13.287	< -1.513
Ni II	1741.6	< 13.287		
Ni II	1751.9	< 13.494		
Zn II	2026.1	< 12.053	< 12.053	< -1.167




FIG. 56.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.224$ toward PSS1506+5220. The vertical line at $v = 0$ corresponds to $z = 3.22442$.




FIG. 57.— Ly α profile of the damped Ly α system at $z = 3.695$ toward PSS1723+2243. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.50^{+0.15}_{-0.15}$. The continuum is difficult to predict for the region surrounding the DLA because it includes the OVI emission peak of the quasar. Furthermore, the profile is best fit by two components. The individual $N(\text{HI})$ values are not well determined but we feel the total column density is accurate to 0.15 dex.




FIG. 58.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.695$ toward PSS1723+2243. The vertical line at $v = 0$ corresponds to $z = 3.69474$.




FIG. 59.— Ly α profile of the damped Ly α system at $z = 3.316$ toward PSS2155+1358. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.55^{+0.15}_{-0.15}$. One notes the presence of several coincident Ly α clouds which absorb the core and red wing of the damped Ly α profile. In this case, we have fit the trace of flux at $\lambda \approx 5258\text{\AA}$ and the few regions of the blue wing which appear uncontaminated. The fit is further complicated by the uncertainty of the continuum and we estimate an error of 0.15 dex to our $N(\text{HI})$ measurement.




FIG. 60.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.316$ toward PSS2155+1358. The vertical line at $v = 0$ corresponds to $z = 3.31625$.

4.24. PSS2155+1358, $z = 3.316$

This PSS quasar shows a damped Ly α system with a complicated Ly α profile (Figure 59). We note that the quasar was observed in poor conditions and has poor S/N for its total exposure time. There is severe blending in the core and red wing of the profile from coincident Ly α clouds which limits the precision of the $N(\text{HI})$ measurement. Our adopted value $N(\text{HI}) = 10^{20.55} \text{ cm}^{-2}$ is significantly lower than the value reported by Peroux et al. (2001). We suspect the difference can be attributed to their inability to identify the coincident Ly α features in their spectra. Figure 60 shows the few metal-line profiles observed for this DLA and Table 33 lists the column density measurements.

4.25. Q2223+20, $z = 3.119$

TABLE 34
IONIC COLUMN DENSITIES: Q2223+20, $z = 3.119$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	13.572 ± 0.024		
C IV	1550.8	13.678 ± 0.034		
N I	1134.4	< 13.619	< 13.619	< -2.611
O I	1302.2	> 14.842	> 14.842	> -2.198
Al II	1670.8	< 12.329	< 12.329	< -2.461
Si II	1260.4	> 13.787	13.640 ± 0.036	-2.220 ± 0.106
Si II	1526.7	13.640 ± 0.036		
Si IV	1393.8	13.236 ± 0.019		
Si IV	1402.8	13.124 ± 0.043		
Fe II	1608.5	13.376 ± 0.090	13.324 ± 0.061	-2.476 ± 0.117
Fe II	2344.2	13.292 ± 0.081		
Fe II	2382.8	> 13.438		
Fe III	1122.5	< 13.600		




FIG. 61.— Ly α profile of the damped Ly α system at $z = 3.119$ toward Q2223+20. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.30^{+0.10}_{-0.10}$. This fit was straightforward except for the determination of the quasar continuum. We note that this DLA only just meets the DLA criterion of $2 \times 10^{20} \text{ cm}^{-2}$ and warn that future observations may remove it from the statistical sample.

This radio-loud quasar was discovered as part of the Third MIT-Green Bank 5GHz Survey (Griffith et al. 1990). Although Storrie-Lombardi and Wolfe (2000) did not identify the $z = 3.119$ absorption system as a DLA, our best fit (Figure 61) just exceeds the DLA criterion of statistical samples. Its metal-line profiles are shown in Figure 62 and Table 34 lists the ionic column densities.

4.26. PSS2241+1352, $z = 4.282$

This damped Ly α system (Peroux et al. 2001) has one of the highest redshifts in our sample. While the determination of its $N(\text{HI})$ value is complicated by the Ly α emission peak (Figure 63), its HI value is constrained by the higher order Lyman lines and our best fit value is $N(\text{HI}) = 10^{21.15} \text{ cm}^{-2}$. At present, this is the largest HI value for a damped system at $z > 4$. The metal-line profiles for the DLA are shown in Figure 64 which includes




FIG. 62.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.119$ toward Q2223+20. The vertical line at $v = 0$ corresponds to $z = 3.11921$.

a probable detection of C II* 1335. The column densities are given by Table 35.

4.27. PSS2323+2758, $z = 3.684$

This PSS quasar exhibits a damped system at $z = 3.684$ with a considerable H I column density $N(\text{H I}) \approx 10^{21} \text{ cm}^{-2}$ (Figure 65). Furthermore, it has one of the lowest metallicities of any DLA, with an Fe abundance below 10^{-3} . The few valuable metal-line profiles are shown in Figure 66 and Table 36 provides the ionic column densities. In passing, we note that a measurement of the O I 1039 transition reveals $N(\text{O}^+) > 15.6$ if one assumes that this profile is unblended. We expect that this is a false assumption but suggest the system warrants further analysis.

4.28. FJ2334-09, $z = 3.057$

TABLE 35
IONIC COLUMN DENSITIES: PSS2241+1352, $z = 4.282$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 15.015	> 15.015	> -2.725
C II*	1335.7	13.467 ± 0.086		
C IV	1550.8	14.055 ± 0.050		
N I	1134.4	< 14.651	> 14.808	> -2.272
N I	1200.2	> 14.809		
Al II	1670.8	> 13.520	> 13.519	> -2.121
Si II	1260.4	> 14.162	> 15.011	> -1.699
Si II	1304.4	> 15.011		
Si II	1526.7	> 14.727		
Si II	1808.0	< 15.103		
Si IV	1393.8	13.376 ± 0.054		
S II	1259.5	14.575 ± 0.033	14.575 ± 0.033	-1.775 ± 0.105
Fe II	1608.5	> 14.646	> 14.646	> -2.004
Fe II	1611.2	< 14.864		
Ni II	1317.2	< 13.223	< 13.223	< -2.177
Ni II	1709.6	< 13.588		
Ni II	1741.5	< 13.507		




FIG. 63.— Ly α profile of the damped Ly α system at $z = 4.282$ toward PSS2241+1352. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{H I}) = 21.15^{+0.10}_{-0.10}$. Although the quasar continuum (dotted line) for this portion of the spectrum is complicated, the core is well isolated from contamination and we are reasonably confident in the best fit value.

TABLE 36
IONIC COLUMN DENSITIES: PSS2323+2758, $z = 3.684$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	13.417 ± 0.066		
O I	1039.2	> 15.607	> 15.607	> -2.083
O I	1302.2	> 14.824		
Al II	1670.8	12.354 ± 0.034	12.354 ± 0.033	-3.086 ± 0.105
Al III	1854.7	12.500 ± 0.127		
Si II	1526.7	13.917 ± 0.030	13.917 ± 0.030	-2.593 ± 0.104
Si II	1808.0	< 14.667		
Si IV	1393.8	< 12.376		
Fe II	1608.5	13.321 ± 0.128	13.321 ± 0.128	-3.129 ± 0.162




FIG. 64.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 4.282$ toward PSS2241+1352. The vertical line at $v = 0$ corresponds to $z = 4.28239$.

This FIRST quasar (White et al. 2000) exhibits a modest damped Ly α system at $z = 3.057$ (Figure 67). The continuum varies significantly across the Ly α profile, but the core constrains the H I column density to reasonable precision. We present the metal-line profiles in Figure 68 and the ionic column densities in Table 37.

4.29. *Q2342+34*, $z = 2.908$

This quasar is another member of the Third MIT-Green Bank 5GHz Survey (Griffith et al. 1990) while the damped Ly α system was identified by White, Kinney, & Becker (1993). The Ly α profile of this damped system (Figure 69 is blended with a neighboring sub-DLA and the red wing is dominated by the Ly α emission peak. Nevertheless, the observations nicely constrain the $N(\text{H I})$ value. Figure 70 plots the comprehensive set of velocity profiles of this DLA and Table 38 gives the ionic column densities.




FIG. 65.— Ly α profile of the damped Ly α system at $z = 3.684$ toward PSS2323+2758. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{H I}) = 20.95^{+0.10}_{-0.10}$. This profile is reasonably well fit even though the blue wing is largely unconstrained by the observations.

TABLE 37

IONIC COLUMN DENSITIES: FJ2334-09, $z = 3.057$

Ion	λ	AODM	N_{adopt}	[X/H]
C II	1334.5	> 15.120	> 15.120	> -1.920
C II*	1335.7	< 12.934		
C IV	1548.2	> 14.389		
C IV	1550.8	> 14.485		
O I	1302.2	> 15.497	> 15.497	> -1.693
Al II	1670.8	> 13.317	> 13.317	> -1.623
Al III	1854.7	12.683 ± 0.047		
Al III	1862.8	12.630 ± 0.082		
Si II	1526.7	> 14.560	14.860 ± 0.070	-1.150 ± 0.122
Si II	1808.0	14.860 ± 0.070		
Si IV	1393.8	> 13.913		
Si IV	1402.8	> 13.924		
Fe II	1608.5	14.334 ± 0.015	14.323 ± 0.014	-1.627 ± 0.101
Fe II	1611.2	< 14.893		
Fe II	2374.5	14.258 ± 0.041		
Ni II	1317.2	< 13.568	< 13.125	< -1.575
Ni II	1741.6	< 13.125		
Zn II	2026.1	< 12.168	< 12.167	< -0.953




FIG. 66.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.684$ toward PSS2323+2758. The vertical line at $v = 0$ corresponds to $z = 3.6845$.




FIG. 67.— Ly α profile of the damped Ly α system at $z = 3.057$ toward FJ2334-09. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 20.45^{+0.10}_{-0.10}$. The continuum varies significantly across the Ly α profile, but the core constrains the HI column density to reasonable precision.

TABLE 38
IONIC COLUMN DENSITIES: Q2342+34, $z = 2.908$

Ion	λ	AODM	N_{adopt}	[X/H]
C I	1656.9	< 12.909		
C II	1334.5	> 15.188	> 15.188	> -2.502
C IV	1548.2	> 14.403		
C IV	1550.8	14.403 ± 0.016		
N I	1135.0	14.887 ± 0.045	14.887 ± 0.045	-2.143 ± 0.110
O I	1302.2	> 15.526	> 15.526	> -2.314
O I	1355.6	< 18.223		
Mg II	1240.4	< 15.612	< 15.612	< -1.068
Al II	1670.8	> 13.615	> 13.615	> -1.975
Al III	1854.7	13.112 ± 0.035		
Si II	1304.4	> 15.047	15.467 ± 0.033	-1.193 ± 0.105
Si II	1526.7	> 14.845		
Si II	1808.0	15.467 ± 0.033		
Si IV	1393.8	> 13.999		
Si IV	1402.8	14.004 ± 0.022		
S II	1259.5	> 15.090	> 15.090	> -1.210
Cr II	2056.3	13.361 ± 0.074	13.361 ± 0.074	-1.409 ± 0.124
Fe II	1122.0	14.982 ± 0.064	14.982 ± 0.064	-1.618 ± 0.119
Fe II	1608.5	> 14.896		
Fe II	1611.2	< 14.942		
Fe II	2249.9	< 15.219		
Fe II	2344.2	> 14.672		
Fe III	1122.5	< 13.940		
Ni II	1709.6	13.852 ± 0.085	13.827 ± 0.045	-1.523 ± 0.110
Ni II	1741.6	13.771 ± 0.069		
Ni II	1751.9	13.922 ± 0.078		
Zn II	2026.1	12.501 ± 0.108	12.501 ± 0.108	-1.269 ± 0.147




FIG. 68.— Velocity plot of the metal-line transitions for the damped $\text{Ly}\alpha$ system at $z = 3.057$ toward FJ2334-09. The vertical line at $v = 0$ corresponds to $z = 3.057195$.




FIG. 69.— $\text{Ly}\alpha$ profile of the damped $\text{Ly}\alpha$ system at $z = 2.908$ toward Q2342+34. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.10^{+0.10}_{-0.10}$. The solution for this $\text{Ly}\alpha$ profile is complicated by a neighboring sub-DLA and the $\text{Ly}\alpha$ emission peak but the blue wing and the flux at $\lambda \approx 4780\text{\AA}$ do nicely constrain the $N(\text{HI})$ value. Note the 1 pixel spike at $\lambda \approx 4750\text{\AA}$ is a cosmic ray.




FIG. 70.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 2.908$ toward Q2342+34. The vertical line at $v = 0$ corresponds to $z = 2.90823$.

Fig 70 – cont

TABLE 39

IONIC COLUMN DENSITIES: PSS2344+0342, $z = 3.219$

Ion	λ	AODM	N_{adopt}	[X/H]
C IV	1548.2	> 14.568		
C IV	1550.8	> 14.726		
N I	1135.0	> 14.648	> 14.648	> -2.632
Al II	1670.8	> 13.535	> 13.535	> -2.305
Al III	1854.7	12.845 ± 0.046		
Si II	1526.7	> 14.866	> 14.865	> -2.045
Cr II	2056.3	< 13.112	13.306 ± 0.088	-1.714 ± 0.112
Cr II	2062.2	13.306 ± 0.088		
Fe II	1608.5	> 14.707	15.234 ± 0.075	-1.616 ± 0.103
Fe II	1611.2	15.234 ± 0.075		
Fe II	2374.5	> 14.517		
Ni II	1709.6	13.546 ± 0.144	13.545 ± 0.144	-2.055 ± 0.160
Ni II	1751.9	< 13.599		
Zn II	2026.1	< 12.267	< 12.267	< -1.753
Zn II	2062.7	< 12.601		




FIG. 71.— Ly α profile of the damped Ly α system at $z = 3.219$ toward PSS2344+0342. The overplotted solid line and accompanying dash-dot lines trace the best fit solution and the estimated bounds corresponding to $\log N(\text{HI}) = 21.35^{+0.07}_{-0.07}$. Because of the well behaved continuum and a relatively clean profile, we assess an uncertainty of only 0.07 dex to this best fit value.

4.30. PSS2344+0342, $z = 3.219$

This damped Ly α system was discovered by Peroux et al. (2001) toward the PSS quasar PSS2344+0342 (Djorgovski 2000). Its Ly α profile is displayed in Figure 71 which is somewhat blended with coincident Ly α lines. The continuum is well behaved across the profile, though, and the $N(\text{HI})$ value is tightly constrained ($10^{21.35 \pm 0.07} \text{ cm}^{-2}$). This HI value is considerable; this DLA has one of the largest column densities at $z > 3$. Note that our value is considerably larger (0.45 dex) than the one reported by Peroux et al. (2001). Given the quality of our fit, we suspect a systematic error in their continuum placement. We also note that Peroux et al. (2001) reported a second DLA along this sightline at $z = 2.68$. Unfortunately, this profile lands at the defect in the ESI CCD which prevents




FIG. 72.— Velocity plot of the metal-line transitions for the damped Ly α system at $z = 3.219$ toward PSS2344+0342. The vertical line at $v = 0$ corresponds to $z = 3.2194$.

us from confirming its existence. We do worry, however, that the higher order Lyman series and Lyman limit of a $z = 3.93$ system do cut into the spectrum at the position where they reported the Ly α profile. Also, we have been unable to identify and Fe II, Zn II, or Cr II transitions with the claimed DLA. For these reasons, we suspect the DLA was a mistaken identification.

An extensive set of metal-line profiles of the $z = 3.2194$ DLA are presented in Figure 72 and the column densities are listed in Table 39. One notes that the system appears to exhibit $[\text{Zn}/\text{Fe}] < 0$, implying a very low dust content.

5. SUMMARY

Tables 40 and 41 present a summary of the absolute and relative abundances of the 37 damped Ly α systems in our complete database. In Table 41, where we present abundances relative to Fe, we have considered Ni or Al as

TABLE 40
ABUNDANCE SUMMARY

Name	z_{abs}	$N(\text{HI})$	[C/H]	[N/H]	[O/H]	[Al/H]	[Si/H]	[S/H]	[Ti/H]	[Cr/H]	[Fe/H]	[Ni/H]	[Zn/H]
SDSS0127-00	3.727	21.15	> -3.068	-3.052	> -2.705	> -2.897	> -2.404					< -1.709	
PSS0133+0400	3.693	20.70	> -3.130			-2.818	> -2.004				-2.689		
PSS0133+0400	3.774	20.55	> -1.831			> -1.248	-0.644		-0.449	< -0.980	> -1.182	-0.822	< -0.121
PSS0134+3317	3.761	20.85				-2.728	> -2.521						
PSS0209+0517	3.667	20.45		< -2.588	> -2.416	-2.420	> -1.903			< -1.041	-2.315	< -1.450	
PSS0209+0517	3.864	20.55	> -3.120		> -2.922	-2.894	-2.647				< -2.710	< -1.677	
BRJ0426-2202	2.983	21.50				> -3.059	< -2.064			< -2.269	-2.848	< -2.377	< -2.005
FJ0747+2739	3.423	20.85	> -2.344			> -2.136	> -1.871				> -1.920	< -1.828	
FJ0747+2739	3.900	20.50	> -2.587		> -2.434	-2.529	-2.029	< -1.339			< -2.201	< -1.644	< -0.772
PSS0808+52	3.113	20.65		< -1.356	> -2.445		-1.611	< -0.929		< -1.585	-1.980	< -1.662	< -1.186
FJ0812+32	2.626	21.35	> -2.800	< 0.326	> -2.602	> -2.181	> -1.128	> -1.112	-1.921	-1.700	-1.753	-1.754	-0.985
Q0930+28	3.235	20.35	> -2.767		> -2.894	-2.471	-2.117				-2.360		
PC0953+47	3.404	21.15				> -2.280	> -2.093			< -1.259	> -2.178	< -1.504	
PC0953+47	3.891	21.20				> -1.859	> -1.599				> -1.712	-1.698	
PC0953+47	4.244	20.90	> -2.771		> -2.478	< -2.517	-2.225	< -1.382			-2.500	< -1.537	
PSS0957+33	3.280	20.45		< -1.936		> -1.812	-1.334	< -1.073		< -1.229	> -1.862	< -1.548	< -0.952
PSS0957+33	4.180	20.65	> -2.441		> -2.418	-2.102	> -1.834				> -2.253	< -1.952	
BQ1021+3001	2.949	20.70	> -2.704		> -2.592	-2.497	-2.170			< -1.537	-2.320	< -1.582	< -1.181
CTQ460	2.777	21.00	> -2.439		> -2.176	-2.065	> -1.590	-1.409			-1.662	-1.816	< -1.311
HS1132+2243	2.783	21.00	> -3.010	-2.911	> -2.751	> -2.623	-2.069	-2.129			-1.836	-2.476	< -2.095
Q1209+0919	2.584	21.40				> -1.822	> -0.970		< -1.688	-1.489	-1.681	< -1.432	-1.088
PSS1248+31	3.696	20.63				-2.343	-1.797			< -1.058	-2.240	< -1.387	< -0.685
PSS1253-0228	2.783	21.85	> -3.428		> -3.243	> -2.941	> -1.813		< -1.948	-1.888	-1.992	-1.930	-1.752
Q1337+11	2.795	20.95	> -3.057	-3.129	> -2.824	-2.596	-1.792	-1.880		< -1.720	-2.388	< -2.001	< -1.419
PKS1354-17	2.780	20.30	> -2.748			-2.395	-1.884				-2.426		
PSS1432+39	3.272	21.25				> -1.991	-1.144			< -1.760	> -1.820	-1.750	< -1.275
Q1502+4837	2.570	20.30	> -1.716		> -1.699	> -1.428	-1.619				-1.649		
PSS1506+5220	3.224	20.67	> -2.050		> -2.753	> -2.657	-2.348			< -1.556	-2.462	< -1.507	< -1.235
PSS1723+2243	3.695	20.50				> -1.426	> -1.199			< -0.956	> -1.425	< -0.800	> -0.662
PSS2155+1358	3.316	20.55				> -1.724	-1.263			< -1.069	> -1.677	< -1.513	< -1.167
Q2223+20	3.119	20.30		< -2.611	> -2.198	< -2.461	-2.220				-2.476		
PSS2241+1352	4.282	21.15	> -2.725	> -2.272		> -2.121	> -1.699	-1.775			> -2.004	< -2.177	
PSS2323+2758	3.684	20.95			> -2.083	-3.086	-2.593				-3.129		
FJ2334-09	3.057	20.45	> -1.920		> -1.693	> -1.623	-1.150				-1.627	< -1.575	< -0.953
Q2342+34	2.908	21.10	> -2.502	-2.143	> -2.314	> -1.975	-1.193	> -1.210		-1.409	-1.618	-1.523	-1.269
PSS2344+0342	3.219	21.35		> -2.632		> -2.305	> -2.045			-1.714	-1.616	-2.055	< -1.753

TABLE 41
RELATIVE ABUNDANCE SUMMARY

Name	z_{abs}	$N(\text{HI})$	[C/Fe]	[N/Fe]	[O/Fe]	[Al/Fe]	[Si/Fe]	[S/Fe]	[Ti/Fe]	[Cr/Fe]	[Ni/Fe]	[Zn/Fe]
SDSS0127-00	3.727	21.15										
PSS0133+0400	3.693	20.70	> -0.441			-0.129	> +0.685					
PSS0133+0400 ^a	3.774	20.55	> -1.009			> -0.426	+0.178		+0.373	< -0.158		< +0.701
PSS0134+3317 ^b	3.761	20.85					> +0.207					
PSS0209+0517	3.667	20.45		< -0.273	> -0.101	-0.105	> +0.412			< +1.274	< +0.865	
PSS0209+0517 ^b	3.864	20.55	> -0.226		> -0.028		+0.247				< +1.217	
BRJ0426-2202	2.983	21.50				> -0.211	< +0.784			< +0.579	< +0.471	< +0.843
FJ0747+2739	3.423	20.85										
FJ0747+2739 ^b	3.900	20.50	> -0.058		> +0.095		+0.500	< +1.190			< +0.885	< +1.757
PSS0808+52	3.113	20.65		< +0.624	> -0.465		+0.369	< +1.051		< +0.395	< +0.318	< +0.794
FJ0812+32	2.626	21.35	> -1.047	< +2.079	> -0.849	> -0.428	> +0.625	> +0.641	-0.168	+0.053	-0.001	+0.768
Q0930+28	3.235	20.35	> -0.407		> -0.534	-0.111	+0.243					
PC0953+47 ^b	3.404	21.15					> +0.187			< +1.021	< +0.776	
PC0953+47 ^a	3.891	21.20				> -0.161	> +0.099					
PC0953+47	4.244	20.90	> -0.271		> +0.022	< -0.017	+0.275	< +1.118			< +0.963	
PSS0957+33	3.280	20.45										
PSS0957+33 ^b	4.180	20.65	> -0.339		> -0.316		> +0.268				< +0.150	
BQ1021+3001	2.949	20.70	> -0.384		> -0.272	-0.177	+0.150			< +0.783	< +0.738	< +1.139
CTQ460	2.777	21.00	> -0.623		> -0.360	-0.249	> +0.226	+0.407		+0.154	+0.153	< +0.505
HS1132+2243	2.783	21.00	> -0.534	-0.435	> -0.275	> -0.147	+0.407	+0.347		+0.640	< +0.381	< +0.792
Q1209+0919	2.584	21.40				> -0.141	> +0.711		< -0.007	+0.192	< +0.249	+0.593
PSS1248+31	3.696	20.63				-0.103	+0.443			< +1.182	< +0.853	< +1.555
PSS1253-0228	2.783	21.85	> -1.436		> -1.251	> -0.949	> +0.179		< +0.044	+0.104	+0.062	+0.240
Q1337+11	2.795	20.95	> -0.669	-0.741	> -0.436	-0.208	+0.596	+0.508		< +0.668	< +0.387	< +0.969
PKS1354-17	2.780	20.30	> -0.322			+0.031	+0.542					
PSS1432+39 ^a	3.272	21.25				> -0.241	+0.606			< -0.010		< +0.475
Q1502+4837	2.570	20.30	> -0.067		> -0.050	> +0.221	+0.030					
PSS1506+5220	3.224	20.67	> +0.412		> -0.291	> -0.195	+0.114			< +0.906	< +0.955	< +1.227
PSS1723+2243 ^b	3.695	20.50					> +0.227			< +0.470	< +0.626	> +0.764
PSS2155+1358 ^b	3.316	20.55					+0.461			< +0.655	< +0.211	< +0.557
Q2223+20	3.119	20.30		< -0.135	> +0.278	< +0.015	+0.256					
PSS2241+1352	4.282	21.15										
PSS2323+2758	3.684	20.95			> +1.046	+0.043	+0.536					
FJ2334-09	3.057	20.45	> -0.293		> -0.066	> +0.004	+0.477				< +0.052	< +0.674
Q2342+34	2.908	21.10	> -0.884	-0.525	> -0.696	> -0.357	+0.425	> +0.408		+0.209	+0.095	+0.349
PSS2344+0342	3.219	21.35		> -1.016		> -0.689	> -0.429			-0.098	-0.439	< -0.137

^aNi is serving as a proxy for Fe^bAl is serving as a proxy for Fe

a proxy for Fe in a few cases as noted.

We have presented ionic column density measurements for our complete sample of damped Ly α systems. We have measured metal-line column densities with the apparent optical depth method and $N(\text{HI})$ values through qualitative fits to the Ly α profiles. Therefore, all of the data has been reduced and analysed with an identical approach. We have used the most up to date atomic data and will continue to update the database as new information becomes available. Visit <http://www.ucolick.org/~xavier/ESI/index.html> for tables, figures and updated measurements as well as <http://kingpin.ucsd.edu/~hiresdla/> for the HIRES abundance database. Future papers will present new scientific results based on the ESI database. It is our intention to release all of the reduced spectra within 3 years of publication. Monitor the ESI data website for details.

The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain. We also acknowledge the Keck support staff for their efforts in performing these observations. We thank the referee Sara Ellison for her extensive and extremely valuable comments and criticism. E.G. is supported by Fundación Andes and by an NSF Astronomy and Astrophysics Postdoctoral Fellowship under award AST-0201667.

REFERENCES

- Bergeson, S.D. & Lawler, J.E. 1993, *ApJ*, 408, 382
 Bergeson, S.D. & Lawler, J.E. 1993, *ApJ*, 414, L137
 Fedchak, J. A. & Lawler, J. E. 1999, *ApJ*, 523, 734
 Bergeson, S.D., Mullman, K.L., & Lawler, J.E. 1993, *ApJ*, 435, L157
 Bergeson, S.D., Mullman, K.L., Wickliffe, M.W., Lawler, J.E., Litzen, U., and Johansson, S. 1996, *ApJ* 464, 1044
 Dessauges-Zavadsky, M., D’Odorico, S., McMahon, R.G., Molaro, P., Ledoux, C., Péroux, C., & Storrie-Lombardi, L.J. 2001, *A&A*, 370, 426
 Djorgovski, S.G., Gal, R.R., Odewahn, S.C., de Carvalho, R.R., Brunner, R., Longo, G., & Scaramella, R. 1998, in “Wide Field Surveys in Cosmology”, eds. S. Colombi & Y. Mellier, (astro-ph/9809187)
 Djorgovski, S.G., priv. comm., (<http://astro.caltech.edu/~george/z4.qsos>)
 Ellison, S.L., Ryan, S., & Prochaska, J.X. 2001, *MNRAS*, 326, 628
 Ellison, S.L., Yan, L., Hook, I.M., Pettini, M., Wall, J.V., & Shaver, P. 2002, *A&A*, 383, 91
 Fan, X., SDSS collaboration, 2001, *AJ*, 121, 31
 Fedchak, J. A., Wiese, L. M., & Lawler, J. E. 2000, *ApJ*, 538, 773
 Grevesse, N., Noels, A., & Sauval, A.J. 1996, In: *Cosmic Abundances*, S. Holt and G. Sonneborn (eds.), ASPCS, V. 99, (BookCrafters: San Francisco), p. 117
 Gregg, M.D., Becker, R.H., White, R.L., Helfand, D.J., McMahon, R.G., & Hook, I.M. 1996, *AJ*, 112, 407
 Griffith, M., Langston, G., Heflin, M., Conner, S., Lehar, J & Burke, B. 1990, *ApJS*, 74, 129.
 Hagen, H.J., Engels, D., & Reimers, D. 1999, *A&AS*, 134, 483
 Holweger, H. 2001, in *Solar and Galactic Composition*, ed. R.F. Wimmer-Schweingruber, (Berlin: Springer), 23
 Howk, J.C., Sembach, K.R., Roth, K.C., & Kruk, J.W. 2000, *ApJ*, 544, 867
 Lopez, S., Maza, J., Masegosa, J., & Marquez, I. 2001, *A&A*, 366, 387
 Lopez, S., Reimers, D., D’Odorico, S., & Prochaska, J.X. 2002, *A&A*, 385, 778
 Lu, L., Sargent, W.L.W., & Barlow, T.A. 1998, 115, 55
 Maza, J., Wischnjewsky, J., Antezana, R. & Gonzalez, L.E. 1995, *RMx&A*, 31, 119
 Molaro, P., Bonifacio, P., Centurión, M., D’Odorico, S., Vladilo, G., Santin, P., & Di Marcantonio, P. 2000, *ApJ*, 541, 54
 Møller, P., Warren, S.J., & Fynbo, J.U. 1998, *A&A*, 330, 19
 Morton, D.C. 1991, *ApJS*, 77, 119
 Morton, D. 2003, in prep
 Péroux, C., Storrie-Lombardi, L.J., McMahon, R.G., Irwin, M., & Hook, I.M. 2001, *AJ*, 121, 1799
 Petitjean, P., Srianand, R., & Ledoux, C. 2000, *A&A*, 364, 26L
 Pettini, M., Ellison, S.L., Bergeron, J., & Petitjean, P. 2002, *A&A*, 391, 21
 Prochaska, J.X., Gawiser, E., & Wolfe, A.M. 2001, *ApJ*, 552, 99 (PGW01)
 Prochaska, J.X., Howk, J.C., & Wolfe, A.M. 2003, *Nature*, in press
 Prochaska, J. X. & Wolfe, A. M. 1999, *ApJS*, 121, 369
 Prochaska, J.X. & Wolfe, A.M., 2000, *ApJ*, 533, L5
 Prochaska, J.X. & Wolfe, A.M. 2002, *ApJ*, 566, 68
 Prochaska, J.X., Wolfe, A.M., Tytler, D., Burles, S.M., Cooke, J., Gawiser, E., Kirkman, D., O’Meara, J.M., & Storrie-Lombardi, L. 2001, *ApJS*, 137, 21
 Prochaska, J.X., Henry, R.B.C., O’Meara, J.M., Tytler, D., Wolfe, A.M., Kirkman, D., Lubin, D., & Suzuki, N. 2002, *PASP*, 114, 933
 Raassen, A.J.J. & Uylings, P.H.M. 1998, *A&A*, 340, 300
 Sargent, W.L.W., Steidel, C.C., & Boksenberg, A. 1989, *ApJS*, 69, 703
 Savage, B. D. and Sembach, K. R. 1991, *ApJ*, 379, 245
 Schechtman, R.M., Povolny, H.S., & Curtis, L.J. 1998, *ApJ*, 504, 921
 Schilizzi, R.T. & McAdam, W.B. 1975, *MmRAS* 79, 1
 Schneider, D.P., Schmidt, M. & Gunn, J.E. 1991, *AJ* 101, 2004.
 Sheinis, A.I., Miller, J., Bigelow, B., Bolte, M., Epps, H., Kibrick, R., Radovan, M., & Sutin, B. 2002, *PASP*, 114, 851
 Snellen, I.A.G., McMahon, R.G., Dennett-Thorpe, J., Jackson, N., Mack, K.-H. & Xanthopoulos, E. 2001, *MNRAS*, 325, 1167
 Storrie-Lombardi, L.J. & Wolfe, A.M. 2000, *ApJ*, 543, 552
 Storrie-Lombardi, L. J., Irwin, M.J., McMahon, R. G., & Hook, I.M. 2001, *MNRAS*, 322, 933
 Tripp, T. M., Lu L., & Savage B.D. 1996, *ApJS*, 102, 239
 Verner, D. A. 1996, *Atomic Data, Nuc. Data Tables*, 64, 1
 Vogt, S.S., Allen, S.L., Bigelow, B.C., Bresee, L., Brown, B., et al. 1994, *SPIE*, 2198, 362
 White, R.L., Kinney, A.L. & Becker, R.H. 1993, *ApJ* 407, 456
 White, R.L., et al. 2000, *ApJS*, 126, 133
 Wiese, L.M., Fedchak, J. A., & Lawler, J. E. 2001, *ApJ*, 547, 1178
 Wolfe, A. M., Prochaska, J.X., & Gawiser, E. 2003, *ApJS*, in press
 Wolfe, A. M., Lanzetta, K. M., Foltz, C. B., and Chaffee, F. H. 1995, *ApJ*, 454, 698